

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Journal of Econometrics

journal homepage: www.elsevier.com/locate/jeconom

Inference in continuous systems with mildly explosive regressors[☆]

Ye Chen^a, Peter C.B. Phillips^{b,c,d,e,*}, Jun Yu^d

^a Capital University of Economics and Business, China

^b Yale University, United States

^c University of Auckland, New Zealand

^d Singapore Management University, Singapore

^e University of Southampton, United Kingdom

ARTICLE INFO

Article history:

Available online xxx

JEL classification:

C12

C13

C58

Keywords:

Cointegrated system

Explosive process

Moderate deviations from unity

Double asymptotics

Real estate market

ABSTRACT

New limit theory is developed for co-moving systems with explosive processes, connecting continuous and discrete time formulations. The theory uses double asymptotics with infill (as the sampling interval tends to zero) and large time span asymptotics. The limit theory explicitly involves initial conditions, allows for drift in the system, is provided for single and multiple explosive regressors, and is feasible to implement in practice. Simulations show that double asymptotics deliver a good approximation to the finite sample distribution, with both finite sample and asymptotic distributions showing sensitivity to initial conditions. The methods are implemented in the US real estate market for an empirical application, illustrating the usefulness of double asymptotics in practical work.

© 2017 Elsevier B.V. All rights reserved.

1. Introduction

The recent global financial crisis has motivated econometricians to study potentially explosive behavior in financial time series and develop technologies for the detection of bubbles in financial markets. For example, Phillips et al. (2011) and Phillips et al. (2015a, b) use mildly explosive representations to capture market exuberance in financial time series and recursive regressions to provide dating algorithms. Both these studies use machinery that draws on work of Phillips and Magdalinos (2007, PM hereafter) on mildly explosive processes and the limit theory associated with these processes, which have a growing number of applications in economics and finance. Other recent research has focussed on mechanisms for generating financial bubbles rather than reduced form methods. Among his many wide-ranging contributions to econometrics and finance, Christian Gouriéroux has recently explored new ways of generating explosive bubbles via non-causal forward-looking processes (Gouriéroux and Zakoian, 2017).

Long run equilibrium relationships among nonstationary variables are often modeled in terms of cointegrated systems. In a

typical cointegrated system variables are assumed to be integrated $I(1)$ processes and the model is formulated in discrete time. However, financial applications often use continuous time representations, given the presence of high frequency observations, making these representations popular in empirical work. Phillips (1991) showed how to formulate a cointegrated system in continuous time and proposed an inferential procedure for such systems based on frequency domain techniques. That work maintained the usual $I(1)$ process assumption, thereby excluding episodes of exuberance in the data.

Extending the framework of co-movement in data to mildly explosive variables, Magdalinos and Phillips (2009, MP hereafter) developed a generalized cointegrated system with multiple variables that may be mildly explosive, leading to mixed normal limit theory and mildly explosive rates of convergence, just as in the univariate autoregression studied in PM. Like autoregressive roots that are local to unity, mildly explosive roots depend on the sample size but deliver parameterizations that lie in a wider vicinity of unity. The limit theory in such systems is independent of the initial condition when, as is often the case, the initialization is assumed to be asymptotically negligible relative to the order of the sample observations. Other cases, where the initialization is non-negligible and may figure in the limit theory in various ways are considered in other work (Andrews and Guggenberger, 2012; Phillips and Magdalinos, 2009).

In a recent study, Wang and Yu (2016, WY hereafter) developed a double asymptotic theory for an explosive continuous time

[☆] Phillips acknowledges support from the NSF under Grant No. SES 12-58258. Yu acknowledges financial support by the Singapore Ministry of Education (MOE) Academic Research Fund Tier 3 grant MOE2013-T3-1-009. We thank two referees, the editors, and seminar participants at Peking University for helpful comments on earlier versions of the paper.

* Corresponding author at: Yale University, United States.
E-mail address: peter.phillips@yale.edu (P.C.B. Phillips).

model, where the sampling interval passes to zero and the time span passes to infinity. In this double asymptotic setting, the explosive continuous time model implies mildly explosive behavior in discrete time but with an autoregressive parameter that depends on the sampling frequency, not the sample size, by virtue of the discrete time solution of the continuous system. In empirical work the value of the autoregressive coefficient is also often taken to depend on the frequency of observation. This is because the use of higher frequency data typically leads to a more persistent autoregressive coefficient estimate and expectations do not change over short time horizons as much as they do over long horizons. For these reasons dependence of the autoregressive parameter on sampling frequency often provides greater realism in empirical work where it is necessary to model near unit root phenomena in discrete time. The limit theory in WY contains a term that explicitly depends on initial conditions, thereby differing from the (large span) limit theory in PM. This difference arises from the different order of magnitude implied for the initial conditions in the two approaches. Simulations in WY reveal that double asymptotics involving initial condition dependencies typically outperform in finite samples the asymptotics that are free of the initial condition. The changes in the limit theory induced by these initial condition dependencies are sufficient to materially change conclusions in empirical work.

This paper extends work by Phillips (1991) on continuous system cointegration by developing asymptotics for continuous models where the variables are mildly explosive. The model differs from MP's mildly explosive system in three ways. First, our model is formulated and parameterized in a way that relates to a continuous time system whereas MP use a discrete time specification. This difference is important because the implied (discrete time) autoregressive parameter of the continuous system depends on the known sampling frequency, not on the sample size in terms of an unknown localizing coefficient. Pivotal limit theory is therefore possible in the continuous time formulation. Second, the initial conditions in the two models are different. Third, the continuous time model allows for a drift in the regressor, which affects the limit theory. In developing double asymptotics, we utilize the limit theory of MP while adjusting for the initial condition, the drift, and the autoregressive specification, all of which affect the resulting limit distribution.

There are good reasons for extending discrete time cointegrated systems to continuous time. Continuous time models now enjoy a wide range of empirical applications both in macroeconomics and financial economics. They provide for discrete sampling at any frequency, including intermittent random sampling, and they allow for convenient handling of both stock variables and flow variables, the latter by simple time aggregation. Importantly in the present setting, the use of a continuous time framework readily accommodates initial condition and drift effects, with a limit theory that is easy to implement in practice with no nuisance parameters. In particular, the limit theory in the continuous system here depends on a persistence parameter (κ) which is consistently estimable. By contrast, discrete time models with local to unity and mildly integrated or mildly explosive autoregressive parameters typically involve localizing coefficients that enter the limit theory as nuisance parameters and are not generally consistently estimable, thereby complicating inference.

The paper is organized as follows. Section 2 introduces the model and gives our main results, providing connections between the continuous time framework considered here and the discrete time cointegrated systems in MP. The limit theory of MP is modified to allow for a discrete time model with initial condition and drift induced by the continuous system, which assists in delivering double asymptotics for the least squares estimator in the continuous system. Section 3 extends the limit results to the multivariate setting. Section 4 reports simulations studying the finite

sample performance of the methods. An empirical application of the methodology to US real estate data is given in Section 4. Section 5 concludes. Proofs of Theorem 2.1 and Corollary 2.2 are given in Appendix A. Proofs of Theorem 3.1 and 3.2 are provided in an online technical supplement.

2. Continuous systems with a mildly explosive regressor

We start our investigation with the following scalar continuous time model in two variates $y(t)$ and $x(t)$. Here $x(t)$ follows an Ornstein–Uhlenbeck process and the stochastic process $y(t)$ comoves with $x(t)$ as¹

$$y(t) = \beta x(t) + u_0(t), \quad (2.1)$$

$$\begin{aligned} dx(t) &= \kappa (\mu - x(t)) dt + dB_x(t), \\ x(0) &= x_0 = O_p(1), \quad \kappa < 0, \end{aligned} \quad (2.2)$$

where $u_0(t)$ is Gaussian pure noise – a generalized stochastic process in continuous time (see Hannan, 1970 and Phillips, 1991). Formally, we may write $u_0(t) = DB_0(t)$, where $B_0(t) = \sigma_{00}W_0(t)$, W_0 is standard Brownian motion, and $D = d/dt$ is the mean square differential operator. While $u_0(t)$ is unrealizable as a covariance stationary stochastic process in continuous time, the corresponding discrete time process is realizable and takes the form of a pure noise process of independent identically distributed (*iid*) $N(0, \sigma_{00}^2)$ errors. This formulation is convenient in that it corresponds to the discrete time system (2.3) that follows and has been extensively used in modeling microstructure noise effects in the measurement of efficient financial asset prices, as discussed below.² We write $B_x(t) = \sigma_{xx}W_x(t)$ where W_x is a standard Brownian motion that may be correlated with W_0 .

The parameter of central interest for inference is the coefficient β which captures the co-movement between $y(t)$ and $x(t)$. The driver process $x(t)$ follows an Ornstein–Uhlenbeck equation with persistence parameter κ . For $\kappa > 0$ the process $x(t)$ is stationary, for $\kappa = 0$ it is Brownian motion, and for $\kappa < 0$ it is explosive. For data over a large time span several different regimes of κ might be contemplated, possibly with break points separating the regimes. The present paper focuses on the explosive case of $\kappa < 0$. The scalar model is important particularly in financial applications and leads to simple results that avoid some of the complications of systems with multiple explosive regressors, which are considered in the next section.

Suppose data are recorded at N equally spaced points, $\{t\}_{t=1}^N$, over a time interval $[0, T]$, with sampling interval h and overall time span T so that $N = T/h$. To develop asymptotics we assume that both $h \rightarrow 0$ and $T \rightarrow \infty$. The exact discrete time representation of (2.1)–(2.2) is (Phillips, 1972)

$$y_{th} = \beta x_{th} + u_{0,th}, \quad (2.3)$$

$$x_{th} = a_h(\kappa) x_{(t-1)h} + g_h + u_{x,th}, \quad x_{0h} = x_0 = O_p(1), \quad (2.4)$$

where

$$\begin{aligned} a_h(\kappa) &= \exp(-\kappa h), \\ g_h &= \mu(1 - e^{-\kappa h}), \end{aligned}$$

¹ Alternate specifications are possible. For example, as discussed in Remark 12 we might set $u_0(t) = a_s \cdot 0$, which corresponds to the limit form of a discrete time cointegrated system. A more general model specification might allow for a realizable covariance stationary error in (2.1). Extending the double limit theory developed in the present paper to cover the case of dependent errors is beyond the scope of the present work and will be explored later.

² In such models, the standard formulation of an efficient price subject to unobserved microstructure noise leads to a continuous system involving generalized stochastic processes (see the discussion in Phillips and Yu, (2006).

$$u_{x,th} = \sigma_{xx} \int_{(t-1)h}^{th} e^{-\kappa(th-s)} dB_x(s) \stackrel{d}{=} N\left(0, \frac{\sigma_{xx}^2}{2\kappa} (1 - e^{-2\kappa h})\right),$$

$$u_{0,th} \stackrel{d}{=} N(0, \sigma_{00}^2).$$

Unlike the variance of the discrete time error $u_{x,th}$ for the process x_{th} , the variance of $u_{0,th}$ does not depend on the sampling interval h and, as implied in the above discussion, $\{u_{0,th}\}_{t=1}^N$ therefore corresponds to an iid Gaussian sequence in discrete time that is associated with the generalized stochastic process $u_0(t)$ in continuous time. The autoregressive parameter $a_h(\kappa)$ depends directly on the sampling frequency h . Indirectly, h and $a_h(\kappa)$ are both related to the sample size N . When T is fixed, $h = T/N = O(N^{-1}) \rightarrow 0$, and when $T \rightarrow \infty$, $h = O(T/N) \rightarrow 0$. Gaussianity follows from the Brownian motion driver processes in (2.1)–(2.2). The standard error of $u_{x,th}$ is $\lambda_h \sim \sigma_{xx}\sqrt{h} \rightarrow 0$, concordant with the sample path continuity of $x(t)$.

Re-standardizing Eq. (2.4) by λ_h we have

$$y_{th} = \beta x_{th} + u_{0,th}, \tag{2.5}$$

$$\tilde{x}_{th} = a_h(\kappa) \tilde{x}_{(t-1)h} + \tilde{g}_h + \tilde{u}_{x,th}, \quad \tilde{x}_{0h} = x_{0h}/\lambda_h,$$

$$\tilde{u}_{x,th} \stackrel{iid}{\sim} N(0, 1), \tag{2.6}$$

where $\tilde{x}_{th} = x_{th}/\lambda_h$, $\tilde{g}_h = g_h/\lambda_h$ as $h \rightarrow 0$. When $T \rightarrow \infty$ and $h \rightarrow 0$, we have

$$\frac{1}{Nh} = \frac{1}{T} \rightarrow 0, \quad a_h(\kappa) = 1 - \kappa h + O(h^2) \rightarrow 1.$$

Hence, \tilde{x}_{th} in (2.6) is a mildly explosive process as in PM. Furthermore, since $\kappa < 0$, when $h \rightarrow 0$ we have $\tilde{x}_{0h} = x_{0h}/\lambda_h = O_p(h^{-1/2})$, $(a_h(\kappa))^{-N} = o(1/T)$, and $\tilde{g}_h \sim O(\sqrt{h})$. Thus, in the standardized discrete system (2.5)–(2.6) the order of magnitude of the initial condition is $\tilde{x}_{0h} \sim O_p(h^{-1/2})$ while in the original system (2.1)–(2.2) it is $x_0 \sim O_p(1)$. In addition, the order of magnitude of the drift is $O(\sqrt{h})$ in model (2.6) but is $O_p(1)$ in (2.2).

The continuous time system (2.1)–(2.2) is formulated in a manner similar to models used in the market microstructure literature, where the observed transaction price is often assumed to differ from the efficient price by an iid noise component—see Zhang et al. (2005), Ait-Sahalia et al. (2005, AMZ hereafter) and Bandi and Russell (2006). In AMZ, the logarithmic efficient price of a security follows

$$dX(t) = \mu(X(t)) dt + \sigma(X(t)) dW(t), \tag{2.7}$$

where $W(t)$ is a standard Brownian motion, and the logarithmic price is observed with an iid error, namely,

$$\tilde{X}(th) = X(th) + U(th), \tag{2.8}$$

where $U(th)$ is iid noise with mean zero and a finite variance σ_u^2 that is independent of the sampling interval h , and $U(th)$ is independent of $X(th)$. The formulation (2.8) therefore corresponds to (2.3) and is a discrete time version of the continuous process co-movement $\tilde{X}(t) = X(t) + U(t)$ with a generalized stochastic process error $U(t)$. Issues of formulating such systems in continuous time have been discussed in this market microstructure noise literature to which readers are referred (e.g. Hansen and Lunde, 2006; Phillips and Yu, 2006).

The continuous time system here also relates to the model in Phillips (1991), who studied the system

$$y_1(t) = By_2(t) + u_1(t),$$

$$Dy_2(t) = u_2(t),$$

where $D = d/dt$ denotes the mean square differential operator and $u(t) = [u_1(t), u_2(t)]'$ is a stationary time series. In Phillips

(1991), both $y_1(t)$ and $y_2(t)$ are $I(1)$ processes and co-movement is a form of continuous time cointegration. In contrast, our focus of attention in the present paper is co-movement among potentially explosive processes.

The standardized discrete system (2.5)–(2.6) is related to MP. MP analyzed the triangular system

$$y_t = Ax_t + u_{0t}, \tag{2.9}$$

$$x_t = R_N x_{t-1} + u_{xt}, \quad x_0 = o_p(N^{\alpha/2}), \tag{2.10}$$

where $R_N = I_K + \frac{C}{N^\alpha}$, $\alpha \in (0, 1)$, $C = \text{diag}(c_1, \dots, c_K)$, and discrete observations $\{y_t, x_t\}_{t=0}^N$ are available. In this system, A is the matrix of cointegrating (or, more specifically in the development below, co-mildly explosive) coefficients; R_N represents moderate deviations from a unit root in the sense of PM; x_t is a moderately integrated time series as $N^\alpha \rightarrow \infty$ when $N \rightarrow \infty$. If $C > 0$, x_t is a mildly explosive time series. The vector (u_{0t}, u_{xt}) is a sequence of zero mean, weakly dependent linear process errors which satisfy certain standard regularity conditions. The analysis of MP covers both cases $C > 0$ and $C < 0$, our focus here is on asymptotics for the mildly explosive case $C > 0$.

The formulation of moderate deviations from a unit root with the setting $R_N = 1 + c/N^\alpha$, $\alpha \in (0, 1)$, can be justified from the double asymptotic ($h \rightarrow 0$ and $T \rightarrow \infty$) point of view in continuous time systems, as argued in WY. This justification is similar to that in Boswijk (2001) who justified the block-local-to-unity concept introduced by Phillips et al. (2001, PMX hereafter) in continuous time systems. In PMX, the block-local-to-unity is formulated as $1 + c/m$ where the full sample consists of M blocks of m observations so that $N = Mm$. In PMX and Boswijk (2001), two kinds of asymptotics were considered, namely, (1) M is fixed and $m \rightarrow \infty$; (2) $m \rightarrow \infty$ and then $M \rightarrow \infty$. In case (1), the autoregressive coefficient parameter $1 + \frac{c}{m}$ has the same order of magnitude as the standard local-to-unity model. In case (2), the autoregressive coefficient parameter is further away from unity than that implied by the standard local-to-unity model. An interesting example in case (2) that was considered in PMX is $m = N^\gamma$ and $M = N^{1-\gamma}$, $\gamma \in (0, 1)$. In this case, both m and M are determined by N .

To formalize the link of the notations (i.e. m, M, N) in Boswijk (2001) and those in our paper (i.e. h, T, N) we have $h = \frac{1}{m}$, $T = M$. Clearly our double asymptotics ($T \rightarrow \infty$ and $h \rightarrow 0$) correspond to case (2) in PMX and Boswijk (2001). However, we do not choose N as in PMX and Boswijk (2001) because N is determined after h and T are chosen in our setup.

There are some common features in model (2.5)–(2.6) and the MP model (2.9)–(2.10): both systems imply co-movement between y and x , and in both models x_t may be mildly explosive. But there are also important differences between these discrete time systems. First, the moderate deviations from unity in the autoregressive coefficient take different forms: in (2.6) the autoregressive coefficient is a function of the sampling interval h , whereas in (2.10) it is formulated as a function of the overall sample size N . A second difference is that, while in (2.10) the initial condition for x_t is assumed to be $o_p(N^{\alpha/2}) = o_p(h^{-1/2})$, in (2.6) it is $O_p(h^{-1/2})$, which translates to $x_0 = O_p(1)$ in the original continuous time system (2.1)–(2.2). So, the initial condition in (2.6) has the larger order of magnitude $O_p(h^{-1/2})$, which corresponds to a distant past initialization in the terminology of Phillips and Magdalinos (2009), where it is shown that such initializations do affect the limit theory. The third difference in the models occurs in the drift. In (2.10) there is no intercept, and if a constant intercept were present it would typically dominate the asymptotics. By contrast, in (2.6) the

intercept

$$\begin{aligned} \tilde{g}_h &= g_h/\lambda_h = \mu(1 - e^{-\kappa h})/\lambda_h = \frac{\mu(1 - e^{-\kappa h})}{\sigma_{xx}\sqrt{(1 - e^{-2\kappa h})/2c}} \\ &= \frac{\mu\kappa h}{\sigma_{xx}h^{1/2}}\{1 + o(h)\} = O(\sqrt{h}) \end{aligned}$$

is asymptotically negligible as $h \rightarrow 0$, so the intercept does not affect the double asymptotics. The limit theory of MP is readily modified to take into account this new initial condition and drift.

To fix ideas, consider the modified MP model

$$y_t = Ax_t + u_{0t}, \tag{2.11}$$

$$\begin{aligned} x_t &= \mu + R_N x_{t-1} + u_{xt}, x_0 = x_{0N} = O_p(N^{\alpha/2}), \\ \mu &= O_p(N^{-\alpha/2}). \end{aligned} \tag{2.12}$$

Let $\tilde{x}_0 = x_{0N}N^{-\alpha/2} \Rightarrow X^*$ and $\tilde{\mu} = N^{\alpha/2}\mu \Rightarrow \mu^*$. The error $u_t = [u_{0t}, u_{xt}]'$ is an iid sequence with mean zero and covariance $\begin{bmatrix} \sigma_{00}^2 & \sigma_{0x} \\ \sigma_{0x} & \sigma_{xx} \end{bmatrix}$. This model extends (2.9)–(2.10) by allowing for a larger initial condition and a (local-to-zero) drift. The following theorem gives the limit theory for the LS estimator of A in (2.11) for the case of a single scalar regressor x_t .

Theorem 2.1. For the discrete time system (2.11)–(2.12) with $R_N = 1 + \frac{c}{N^\alpha}$, $\alpha \in (0, 1)$, and $c > 0$, when $N \rightarrow \infty$, we have

- (i) $(R_N^N N^\alpha)^{-1} \sum_{t=1}^N x_t u_{0t} \Rightarrow \frac{\sigma_{00}}{2c} U_0 (\sigma_{xx} U_x + (2c)^{1/2} D)$,
- (ii) $(R_N^N N^\alpha)^{-2} \sum_{t=1}^N x_t^2 \Rightarrow \left(\frac{1}{2c}\right)^2 (\sigma_{xx} U_x + (2c)^{1/2} D)^2$,

where $(U_0, U_x) \stackrel{d}{=} N(0, I_2)$, $D = X^* + \frac{\mu^*}{c}$, and so

$$R_N^N N^\alpha (\hat{A} - A) \Rightarrow 2c \frac{\sigma_{00} U_0}{\sigma_{xx} U_x + (2c)^{1/2} D}. \tag{2.13}$$

Remark 1. The limit result in Theorem 2.1 can be extended to the case where the errors are weakly dependent. Let u_{0t} in the discrete time system (2.11)–(2.12) be a sequence of zero mean and weakly dependent errors, satisfying Assumption LP as defined in MP, so that

$$N^{-1/2} \sum_{t=1}^N u_{0t} \Rightarrow N(0, \omega_{00}^2).$$

where the long-run variance ω_{00}^2 can be decomposed as

$$\omega_{00}^2 = \sigma_{00}^2 + 2\lambda_{00}, \tag{2.14}$$

with variance component σ_{00}^2 and one-sided long-run variance

$$\lambda_{00} := \sum_{h=1}^{\infty} E(u_{0t} u_{0t-h}).$$

Using the limit results from Theorem 4.1 in MP and Theorem 2.1, we obtain

$$R_N^N N^\alpha (\hat{A} - A) \Rightarrow 2c \frac{\omega_{00} U_0}{\sigma_{xx} U_x + (2c)^{1/2} D}. \tag{2.15}$$

Remark 2. If $\tilde{x}_0 = -\frac{\tilde{\mu}}{c}$, then $D = 0$ and the limit (2.13) is simply

$$R_N^N N^\alpha (\hat{A} - A) \Rightarrow 2c \frac{\sigma_{00} U_0}{\sigma_{xx} U_x} = 2c \frac{\sigma_{00}}{\sigma_{xx}} \mathbb{C}, \tag{2.16}$$

where \mathbb{C} is a standard Cauchy variate. This limit distribution is the same as that given by MP (2009, p. 496) and depends on the localizing coefficient c , although the standardized estimation error satisfies

$$\frac{R_N^N}{R_N^2 - 1} (\hat{A} - A) \Rightarrow \frac{\sigma_{00}}{\sigma_{xx}} \mathbb{C}, \tag{2.17}$$

when $D = 0$ and this limit does not depend on c . In the general case where $D \neq 0$

$$\frac{R_N^N}{R_N^2 - 1} (\hat{A} - A) \Rightarrow \frac{\sigma_{00} U_0}{\sigma_{xx} U_x + (2c)^{1/2} D}. \tag{2.18}$$

Remark 3. The limit distribution of $\hat{\mu}$, the LS estimator of the intercept parameter μ , follows simply as given in Box 1. This result is useful in testing for $\mu = 0$ in the modified MP model.

Remark 4. Self normalized statistics based on \hat{A} have a much simpler limit theory that is convenient for inference. For instance, defining the regression residuals $\hat{u}_{0t} = y_t - \hat{A}x_t$ and noting that the residual variance estimate $s_0^2 = N^{-1} \sum_{t=1}^N \hat{u}_{0t}^2 \xrightarrow{p} \sigma_{00}^2$, it follows immediately from Theorem 2.1 that the usual t statistic for testing $\mathbb{H}_0 : A = A^0$ satisfies

$$\begin{aligned} t_A &= \frac{\hat{A} - A^0}{s_A} = \frac{R_N^N N^\alpha (\hat{A} - A^0)}{\left\{s_0^2 \left(\frac{1}{R_N^2 N^{2\alpha}} \sum_{t=1}^N x_t^2\right)^{-1}\right\}^{1/2}} \\ &\Rightarrow \frac{2c \frac{\sigma_{00} U_0}{\sigma_{xx} U_x + (2c)^{1/2} D}}{\sigma_{00} \left\{\left(\frac{1}{2c}\right)^2 (\sigma_{xx} U_x + (2c)^{1/2} D)^2\right\}^{-1/2}} \\ &= U_0 \stackrel{d}{=} N(0, 1), \end{aligned} \tag{2.19}$$

and standard methods of inference apply.

Remark 5. Let \hat{R}_N be the LS estimator of R_N and $\hat{c} = N^\alpha (\hat{R}_N - 1)$. The limit theory for \hat{R}_N and \hat{c} follows from the proof of Theorem 2.1 and Remark 3. Defining the regression residuals $\hat{u}_{xt} = x_t - \hat{R}_N x_{t-1} - \hat{\mu}$ and noting that the residual variance estimate $s_x^2 = N^{-1} \sum_{t=1}^N \hat{u}_{xt}^2 \xrightarrow{p} \sigma_{xx}^2$, we have the following result for the t statistic for testing $\mathbb{H}_0 : R_N = R_N^0$,

$$\begin{aligned} t_{R_N} &= \frac{\hat{R}_N - R_N^0}{s_{R_N}} \\ &= \frac{R_N^N N^\alpha (\hat{R}_N - R_N^0)}{\left\{s_x^2 \left(\frac{1}{R_N^2 N^{2\alpha}} \sum_{t=1}^N x_{t-1}^2 - \frac{1}{N} \left(\frac{1}{R_N^N N^\alpha} \sum_{t=1}^N x_{t-1}\right)^2\right)^{-1}\right\}^{1/2}} \\ &\Rightarrow \frac{2c \frac{\sigma_{xx} U_x}{\sigma_{xx} U_x + (2c)^{1/2} D}}{\sigma_{xx} \left\{\left(\frac{1}{2c}\right)^2 (\sigma_{xx} U_x + (2c)^{1/2} D)^2\right\}^{-1/2}} = U_x \stackrel{d}{=} N(0, 1). \end{aligned}$$

Similarly, given $R_N = 1 + \frac{c}{N^\alpha}$, we have $s_c = N^\alpha s_{R_N}$ and $\hat{c} - c^0 = (\hat{R}_N - R_N^0) N^\alpha$. Hence, if α is known, the t statistic for testing $\mathbb{H}_0 : c = c^0$ is

$$t_c = \frac{\hat{c} - c^0}{s_c} = (\hat{R}_N - R_N^0) N^\alpha N^{-\alpha} s_{R_N}^{-1} \Rightarrow U_x \stackrel{d}{=} N(0, 1).$$

However, if α is unknown, then both the estimate \hat{c} and the standard error $s_c = N^\alpha s_{R_N}$ are unavailable and inference using this limit theory for \hat{c} is infeasible. As discussed below, this infeasible feature of the discrete time case is quite different in continuous time.

Remark 6. The limit distribution (2.13) is a ratio of two independent Gaussian variates and has heavy tails, just as the Cauchy limit in the special case (2.17) where $D = 0$. Observe that

$$\begin{aligned} R_N^N N^\alpha (\hat{A} - A) &\Rightarrow 2c \frac{\sigma_{00} U_0}{\sigma_{xx} U_x + (2c)^{1/2} D} = b \frac{U_0}{U_x + d}, \\ b &= \frac{2c \sigma_{00}}{\sigma_{xx}}, \quad d = \frac{(2c)^{1/2}}{\sigma_{xx}} D. \end{aligned}$$

$$\begin{aligned} & \sqrt{N}(\hat{\mu} - \mu) \\ &= \sqrt{N} \frac{(R_N^N N^\alpha)^{-2} \sum_{t=1}^N x_{t-1}^2 \left(\frac{1}{N} \sum_{t=1}^N u_{xt} \right) - \frac{1}{N} \left\{ (R_N^N N^\alpha)^{-1} \sum_{t=1}^N x_{t-1} \right\} \left\{ (R_N^N N^\alpha)^{-1} \sum_{t=1}^N x_{t-1} u_{xt} \right\}}{\frac{N}{N} \left\{ (R_N^N N^\alpha)^{-2} \sum_{t=1}^N x_{t-1}^2 \right\} - \frac{1}{N} \left\{ (R_N^N N^\alpha)^{-1} \sum_{t=1}^N x_{t-1} \right\}^2} \\ &= \frac{1}{\sqrt{N}} \sum_{t=1}^N u_{xt} + o_p(1) \Rightarrow N(0, \sigma_{xx}^2). \end{aligned}$$

Box I.

When $D \geq 0$, the density of $U = \frac{U_0}{U_x + d}$ is

$$\begin{aligned} p_U(u) &= \frac{e^{-\frac{1}{2}d^2}}{\pi(1+u^2)} \left[1 + \frac{q}{\varphi(q)} \int_0^q \varphi(y) dy \right], \\ \varphi(y) &= \frac{e^{-y^2/2}}{\sqrt{2\pi}}, \quad q = \frac{d}{\sqrt{1+u^2}}, \end{aligned}$$

(e.g., see Marsaglia, 1965) and has Cauchy-like tails.

We have the following expression for the LS estimator $\hat{\beta}$ of the slope coefficient in the continuous time model (2.1), which is given by

$$\begin{aligned} \hat{\beta} - \beta &= \left(\sum_{t=1}^N x_{th}^2 \right)^{-1} \left(\sum_{t=1}^N x_{th} u_{0,th} \right) \\ &= \frac{1}{\lambda_h} \left(\sum_{t=1}^N \tilde{x}_{th}^2 \right)^{-1} \left(\sum_{t=1}^N \tilde{x}_{th} u_{0,th} \right). \end{aligned} \tag{2.20}$$

The associated limit theory is given in the following corollary.

Corollary 2.2. Let $h = N^{-\alpha}$. For the continuous time system (2.1)–(2.2) with $\kappa < 0$, assume that there exists $\alpha \in (0, 1)$, such that $h^{1-\alpha} T^\alpha \rightarrow 1$ as $T \rightarrow \infty$, we have

$$\frac{a_h^N}{\sqrt{h}} (\hat{\beta} - \beta) \Rightarrow (-2\kappa) \frac{\sigma_{00} U_0}{\sigma_{xx} U_x + (-2\kappa)^{1/2} (x_0 - \mu)}. \tag{2.21}$$

Remark 7. The limit result in Corollary 2.2 can be extended to the case where $u_0(t)$ is weakly dependent in (2.1). Using the limit results from Remark 1, we obtain

$$\frac{a_h^N}{\sqrt{h}} (\hat{\beta} - \beta) \Rightarrow (-2\kappa) \frac{\omega_{00} U_0}{\sigma_{xx} U_x + (-2\kappa)^{1/2} (x_0 - \mu)}. \tag{2.22}$$

Remark 8. After multiplying \sqrt{h} on both sides of (2.13), the limit distribution (2.21) follows directly from (2.13) by replacing σ_{xx} , and D in Theorem 2.1 with 1, and $D^* = \frac{x_0 - \mu}{\sigma_{xx}}$, respectively, giving the stated result.

Remark 9. The continuous time counterpart of N^α is $1/h$ which is known for any given data, so there is no need to estimate the rate parameter α . The continuous time counterpart of c is $-\kappa$ which can be consistently estimated by the least squares method as long as $T \rightarrow \infty$. Analogous to (2.19), self normalized statistics are free of nuisance parameters and hypothesis testing about β can be conducted using the residual variance estimate $s_0^2 = N^{-1} \sum_{t=1}^N \hat{u}_{0,th}^2$, which satisfies $s_0^2 \xrightarrow{p} \sigma_{00}^2$. Corollary 2.2 and results (A.11) and (A.12) in Appendix A then give the following double asymptotics for the usual t statistic for testing $\mathbb{H}_0 : \beta = \beta^0$

$$\begin{aligned} t_\beta &= \frac{\hat{\beta} - \beta^0}{s_\beta} = \frac{(\hat{\beta} - \beta^0) a_h^N / \sqrt{h}}{\left\{ s_0^2 (a_h^{-2N} h \sum_{t=1}^N \tilde{x}_{th}^2 \sigma_{xx}^2 h)^{-1} \right\}^{1/2}} \\ &\Rightarrow \frac{(-2\kappa) \frac{\sigma_{00} U_0}{\sigma_{xx} U_x + (-2\kappa)^{1/2} (x_0 - \mu)}}{\sigma_{00} \left\{ \left(\frac{1}{-2\kappa} \right)^2 (U_x + (-2\kappa)^{1/2} D^*)^2 \sigma_{xx}^2 \right\}^{-1/2}} \\ &= U_0 \stackrel{d}{=} N(0, 1), \end{aligned}$$

which leads to feasible inference concerning the slope coefficient β in continuous time, just as in (2.19) for the coefficient A in the modified MP model.

Remark 10. Following Remark 9, we can obtain the double asymptotic distributions for \hat{a}_h and $\hat{\kappa}$. Defining $s_x^2 = N^{-1} \sum_{t=1}^N \hat{u}_{x,th}^2$, which satisfies $h^{-1} s_x^2 \xrightarrow{p} \sigma_{xx}^2$, the t statistic for \hat{a}_h is as follows:

$$\begin{aligned} t_{a_h} &= \frac{\hat{a}_h - a_h^0}{s_{a_h}} \\ &= \frac{a_h^N / h (\hat{a}_h - a_h^0)}{\left\{ s_x^2 \left(a_h^{-2N} h^2 \sum_{t=1}^N \tilde{x}_{(t-1)h}^2 \sigma_{xx}^2 h - \frac{1}{N} \left(a_h^{-N} h \sum_{t=1}^N \tilde{x}_{(t-1)h} \sigma_{xx} \sqrt{h} \right)^2 \right)^{-1} \right\}^{1/2}} \\ &\Rightarrow \frac{-2\kappa \frac{\sigma_{xx} U_x}{\sigma_{xx} U_x + (-2\kappa)^{1/2} D}}{\sigma_{xx} \left\{ \left(\frac{1}{-2\kappa} \right)^2 (\sigma_{xx} U_x + (-2\kappa)^{1/2} D)^2 \right\}^{-1/2}} = U_x \stackrel{d}{=} N(0, 1). \end{aligned}$$

Similarly, given $a_h = \exp(-kh)$, we have $hs_\kappa = s_{a_h} + o_p(h)$. Following WY, we have

$$\begin{aligned} a_h^N (\hat{\kappa} - \kappa^0) &\Rightarrow 2\kappa \frac{\sigma_{xx} U_x}{\sigma_{xx} U_x + (-2\kappa)^{1/2} D}, \\ \text{and} \\ t_\kappa &= \frac{\hat{\kappa} - \kappa^0}{s_\kappa} \\ &= \frac{a_h^N (\hat{\kappa} - \kappa^0)}{\left\{ s_x^2 \left(\frac{1}{a_h^{2N} / h^2} \sum_{t=1}^N x_{(t-1)h}^2 - \frac{1}{N} \left(\frac{1}{a_h^N / h} \sum_{t=1}^N x_{(t-1)h} \right)^2 \right)^{-1} \right\}^{1/2}} \\ &\Rightarrow \frac{-2\kappa \frac{\sigma_{xx} U_x}{\sigma_{xx} U_x + (-2\kappa)^{1/2} D}}{\sigma_{xx} \left\{ \left(\frac{1}{-2\kappa} \right)^2 (\sigma_{xx} U_x + (-2\kappa)^{1/2} D)^2 \right\}^{-1/2}} = U_x \stackrel{d}{=} N(0, 1). \end{aligned}$$

Clearly, t_κ is a feasible statistic for testing $\mathbb{H}_0 : \kappa = \kappa^0$ in contrast to the discrete time case where the test statistic relies on the unknown rate parameter α .

Remark 11. If $x_0 = \mu$, we have $D^* = 0$ and

$$\frac{a_h^N}{\sqrt{h}} (\hat{\beta} - \beta) \Rightarrow (-2\kappa) \frac{\sigma_{00} U_0}{\sigma_{xx} U_x} = (-2\kappa) \frac{\sigma_{00}}{\sigma_{xx}} C. \tag{2.23}$$

Remark 12. An alternate formulation of (2.1) is to set $u_0(t) \stackrel{a.s.}{=} 0$. This formulation corresponds to a discrete system (2.5) in which $u_{0,th} \stackrel{iid}{\sim} N(0, \sigma_{00}^2 h)$ so that the error variance depends on the sampling interval, just as the error $u_{x,th}$, in (2.4). In this case, in view of the scaling effect in the discrete time error, it follows that there is a faster rate of convergence in estimation of β and the limit distribution of $\hat{\beta}$ is given by

$$\frac{a_h^N}{h} (\hat{\beta} - \beta) \Rightarrow (-2\kappa) \frac{\sigma_{00} U_0}{\sigma_{xx} U_x + (-2\kappa)^{1/2} (x_0 - \mu)}, \quad (2.24)$$

when $h \rightarrow 0$ and $T \rightarrow \infty$. In this alternate model, the relationship between $x(t)$ and $y(t)$ is exact in the limit, analogous to the relationship of limit Brownian motion processes (B_y, B_x) of cointegrated discrete series (y_t, x_t) where $x_t = x_{t-1} + u_{xt}$ and $y_t = \beta x_t + u_{0t}$ with (u_{0t}, u_{xt}) stationary and limiting linear relation $B_y(t) = \beta B_x(t)$.

3. Continuous systems with multiple explosive regressors

This section extends the results above to continuous time systems with more than one mildly explosive regressor. We allow for regressors with multiple forms of explosive behavior using the approach developed in MP for discrete systems. As above, we establish the limit theory for a modified MP model that incorporates an intercept term and allows for a larger initial condition. This theory is applied to the continuous system by assuming $T \rightarrow \infty$ and $h \rightarrow 0$. Following MP, two different cases will be examined which lead to somewhat different limit behaviors: (i) when all the regressors have distinct explosive roots; and (ii) when all the regressors share the same explosive root.

3.1. Limit results in the discrete time framework

We start with the following system with multiple mildly explosive regressors, based on MP,

$$y_t = Ax_t + u_{0t}, \quad (3.1)$$

$$x_t = \mu + R_N x_{t-1} + u_{xt}, \text{ with } x_0 = x_{0N} = O_p(N^{\alpha/2})$$

$$\text{and } \mu = O_p(N^{-\alpha/2}). \quad (3.2)$$

In this case, y_t and x_t are $m \times 1$ and $K \times 1$ vectors respectively, and A is a $m \times K$ matrix of coefficients. In addition, $R_N = I_K + C/N^\alpha$ is a $K \times K$ matrix with $C = \text{diag}(c_1, c_2, \dots, c_K) > 0$. We assume that the errors satisfy

$$u_t = [u'_{0t}, u'_{xt}]' \stackrel{iid}{\sim} (0, \Omega) \text{ with } \Omega = \begin{bmatrix} \Omega_{00} & \Omega_{0x} \\ \Omega_{0x} & \Omega_{xx} \end{bmatrix}.$$

Let the standardized initialization and intercept satisfy $\tilde{x}_0 = x_{0N} N^{-\alpha/2} \Rightarrow X^*$ and $\tilde{\mu} = N^{\alpha/2} \mu \Rightarrow \mu^*$. The model now modifies MP in two ways: (i) the initial value for x is $O_p(N^{\alpha/2})$ which is larger than the $o_p(N^{\alpha/2})$ initialization in MP; (ii) a non-zero drift term of order $O_p(N^{-\alpha/2})$ is included. Following closely the approach of MP, we obtain the limit theory for the LS estimator \hat{A} under two scenarios: (i) where C has distinct diagonal elements, i.e., $c_i \neq c_j$ for $i \neq j$; (ii) where C is a scalar matrix and does not have distinct diagonal elements, so that $c_i = c_j$, for all i, j . In what follows we will frequently use a zero affix to denote the true value of the associated element or matrix.

3.1.1. $c_i \neq c_j$ for $i \neq j$

Theorem 3.1. For the discrete time system (3.1)–(3.2) with $R_N = I_K + C/N^\alpha$, $\alpha \in (0, 1)$, $C = \text{diag}(c_1, c_2, \dots, c_K) > 0$ and $c_i \neq c_j$ for $i \neq j$, when $N \rightarrow \infty$, we have

$$\begin{aligned} & (i) \text{vec} \left(\frac{1}{N^\alpha} \sum_{t=1}^N u_{0t} x'_t R_N^{-N} \right) \Rightarrow \left(\int_0^\infty e^{-pC} (D + \tilde{U}_x) (D + \tilde{U}_x)' e^{-pC} dp \otimes \Omega_{00} \right)^{1/2} W_0, \\ & (ii) \frac{1}{N^{2\alpha}} \sum_{t=1}^N R_N^{-N} x_t x'_t R_N^{-N} = \int_0^\infty e^{-pC} (D + \tilde{U}_x) (D + \tilde{U}_x)' e^{-pC} dp + o_p(1), \\ & (iii) \\ & \text{vec} \{ N^\alpha (\hat{A} - A) R_N^N \} \\ & \Rightarrow \left\{ \left(\int_0^\infty e^{-pC} (D + \tilde{U}_x) (D + \tilde{U}_x)' e^{-pC} dp \right)^{-1/2} \otimes \Omega_{00}^{1/2} \right\} W_0 \\ & \stackrel{d}{=} MN \left(0, \left(\int_0^\infty e^{-pC} (D + \tilde{U}_x) (D + \tilde{U}_x)' e^{-pC} dp \right)^{-1} \otimes \Omega_{00} \right), \end{aligned} \quad (3.3)$$

where $W_0 \stackrel{d}{=} N(0, I_{mk})$, $\tilde{U}_x = \left(\int_0^\infty e^{-pC} \Omega_{xx} e^{-pC} dp \right)^{1/2} U_x$, $U_x \stackrel{d}{=} N(0, I_k)$, $D = X^* + C^{-1} \mu^*$, MN represents a mixed normal distribution.

Remark 13. Again, we can extend the limit result in Theorem 3.1 to the case where u_{0t} is weakly dependent. Let u_{0t} in the discrete time system (3.1)–(3.2) be a sequence of zero mean and weakly dependent errors, satisfying Assumption LP as defined in MP. In particular,

$$N^{-1/2} \sum_{i=1}^N u_{0t} \Rightarrow N(0, \Omega_{00}),$$

with long-run variance matrix Ω_{00} decomposed as

$$\Omega_{00} = \Sigma_{00} + \Lambda_{00} + \Lambda'_{00}. \quad (3.4)$$

where

$$\Sigma = \begin{bmatrix} \Sigma_{00} & \Sigma_{0x} \\ \Sigma_{0x} & \Sigma_{xx} \end{bmatrix},$$

and the one-sided long-run covariance is

$$\Lambda_{00} := \sum_{h=1}^\infty E(u_{0t} u'_{0t-h}).$$

Using the limit results from Theorem 3.1 and Theorem 4.1 in MP, we obtain

$$\begin{aligned} & \text{vec} \{ N^\alpha (\hat{A} - A) R_N^N \} \\ & \Rightarrow \left\{ \left(\int_0^\infty e^{-pC} (D + \tilde{U}_x) (D + \tilde{U}_x)' e^{-pC} dp \right)^{-1/2} \otimes \Omega_{00}^{1/2} \right\} W_0 \\ & \stackrel{d}{=} MN \left(0, \left(\int_0^\infty e^{-pC} (D + \tilde{U}_x) (D + \tilde{U}_x)' e^{-pC} dp \right)^{-1} \otimes \Omega_{00} \right). \end{aligned} \quad (3.5)$$

Remark 14. If $\tilde{x}_0 = -C^{-1} \tilde{\mu}$, then $D = 0$ and the limit (3.3) becomes

$$\begin{aligned} & \text{vec} \{ N^\alpha (\hat{A} - A) R_N^N \} \\ & \Rightarrow MN \left(0, \left(\int_0^\infty e^{-pC} \tilde{U}_x \tilde{U}'_x e^{-pC} dp \right)^{-1} \otimes \Omega_{00} \right). \end{aligned}$$

This limit distribution corresponds to that in Theorem 4.1 of MP (2009, p. 496).

Remark 15. The limit distribution of $\widehat{\mu}$ is

$$\begin{aligned} & \sqrt{N}(\widehat{\mu} - \mu) \\ &= \sqrt{N} \left\{ \begin{aligned} & \frac{1}{N} \sum_{t=1}^N u_{xt} - \frac{1}{N} \left(\frac{1}{N^\alpha} \sum_{t=1}^N u_{xt} x'_{t-1} R_N^{-N} \right) \\ & \left(\frac{1}{N^{2\alpha}} \sum_{t=1}^N R_N^{-N} x_{t-1} x'_{t-1} R_N^{-N} \right) \left(\frac{1}{N^\alpha} \sum_{t=1}^N R_N^{-N} x_{t-1} \right) \end{aligned} \right\} \\ & \left\{ \frac{N}{N} - \frac{1}{N} \left(\frac{1}{N^\alpha} \sum_{t=1}^N x'_{t-1} R_N^{-N} \right) \left(\frac{1}{N^{2\alpha}} \sum_{t=1}^N R_N^{-N} x_{t-1} x'_{t-1} R_N^{-N} \right)^{-1} \right. \\ & \quad \left. \times \left(\frac{1}{N^\alpha} \sum_{t=1}^N R_N^{-N} x_{t-1} \right) \right\}^{-1} \\ &= \frac{1}{\sqrt{N}} \sum_{t=1}^N u_{xt} + o_p(1) \Rightarrow N(0, \Omega_{xx}). \end{aligned}$$

Remark 16. Let A_j and \widehat{A}_j denote the j th $m \times 1$ column of A and \widehat{A} , and x_{jt} the j th element of x_t . Define the equation residuals $\hat{u}_{0t} = y_t - \widehat{A}x_t$, the error variance matrix estimate

$$S_{00} = N^{-1} \sum_{t=1}^N \hat{u}_{0t} \hat{u}'_{0t} \xrightarrow{p} \Omega_{00},$$

and the corresponding estimate of the variance matrix of \widehat{A}_j

$$S_{A_j A_j} = \left(\sum_{t=1}^N x_{jt}^2 \right)^{-1} S_{00}.$$

The limit distribution of \widehat{A}_j is given by

$$\begin{aligned} N^\alpha (\widehat{A}_j - A_j) \rho_j^N &\Rightarrow MN \left(0, \left(\int_0^\infty e^{-2pc_j} (D_j + \widetilde{U}_{x_j})^2 dp \right)^{-1} \Omega_{00} \right) \\ &\stackrel{d}{=} MN \left(0, \frac{2c_j \Omega_{00}}{(D_j + \widetilde{U}_{x_j})^2} \right), \end{aligned}$$

where $\rho_j = 1 + \frac{c_j}{N^\alpha}$, and D_j and \widetilde{U}_{jx} are the j th element of D and \widetilde{U}_x for $j = 1, \dots, K$. Using Theorem 3.1, we obtain the following limit distribution for the Wald statistic for testing $\mathbb{H}_0 : Q_j A_j = Q_j A_j^0 = q_j$, where Q_j is a $g \times m$ restriction matrix of full row rank $g \leq m$ and q_j is a given $g \times 1$ vector,

$$\begin{aligned} W_{A_j} &:= \{Q_j \widehat{A}_j - q_j\}' (Q_j S_{A_j A_j} Q_j')^{-1} \{Q_j \widehat{A}_j - q_j\} \\ &= \{Q_j N^\alpha (\widehat{A}_j - A_j) \rho_j^N\}' \left(\left(N^{-2\alpha} \rho_j^{-2N} \sum_{t=1}^N x_{jt}^2 \right)^{-1} Q_j S_{00} Q_j' \right)^{-1} \\ & \quad \times \{Q_j N^\alpha (\widehat{A}_j - A_j) \rho_j^N\} \Rightarrow \chi_g^2, \end{aligned}$$

where χ_g^2 denotes a chi-squared variate with g degrees of freedom.

Remark 17. Let R_{jN} and \widehat{R}_{jN} denote the j th $K \times 1$ column of R_N and \widehat{R}_N , and define $\widehat{C}_j = (\widehat{R}_j - e_j) N^\alpha$, where e_j is the j th unit vector with unity in the j th position and zeros elsewhere. Setting $\hat{u}_{xt} = x_t - \widehat{R}_N x_{t-1} - \widehat{\mu}$, the residual second moment matrix is

$$S_{xx} = N^{-1} \sum_{t=1}^N \hat{u}_{xt} \hat{u}'_{xt} \xrightarrow{p} \Omega_{xx},$$

and the corresponding estimate of the variance matrix of \widehat{R}_{jN} is

$$S_{R_j R_j} = \left(\sum_{t=1}^N x_{jt-1}^2 - \frac{1}{N} \left(\sum_{t=1}^N x_{jt-1} \right)^2 \right)^{-1} S_{xx}.$$

The Wald statistic for testing the (full rank) restrictions $\mathbb{H}_0 : Q_j R_{jN} = Q_j R_{jN}^0 = q_j$, where Q_j is a $g \times K$ restriction matrix of full row rank $g \leq K$ and q_j is a given $g \times 1$ vector, is

$$W_{R_{jN}} := \{Q_j \widehat{R}_{jN} - q_j\}' (Q_j S_{R_j R_j} Q_j')^{-1} \{Q_j \widehat{R}_{jN} - q_j\} \Rightarrow \chi_g^2,$$

under the null. Similarly, given $R_N = I_K + C/N^\alpha$, we can set $S_{C_j C_j} = N^{2\alpha} S_{R_j R_j}$ where C_j is the j th column of C , $S_{C_j C_j}$ is the covariance matrix of \widehat{C}_j . Further $\widehat{C}_j - C_j^0 = (\widehat{R}_{jN} - R_{jN}^0) N^\alpha$ leads to the following limit theory for \widehat{C}_j

$$\begin{aligned} (\widehat{C}_j - C_j^0) \rho_j^N &\Rightarrow MN \left(0, \left(\int_0^\infty e^{-2pc_j} (D_j + \widetilde{U}_{x_j})^2 dp \right)^{-1} \Omega_{xx} \right) \\ &\stackrel{d}{=} MN \left(0, \frac{2c_j \Omega_{xx}}{(D_j + \widetilde{U}_{x_j})^2} \right). \end{aligned}$$

Hence, if α is known, we have the corresponding feasible Wald statistic for testing the restrictions $\mathbb{H}_0 : Q_j C_j = Q_j C_j^0 = q_j$,

$$W_{C_j} := \{Q_j \widehat{C}_j - q_j\}' (Q_j S_{C_j C_j} Q_j')^{-1} \{Q_j \widehat{C}_j - q_j\} \Rightarrow \chi_g^2,$$

under the null and with full row rank Q_j . If α is unknown, just as in the scalar model, the estimated variance matrix $S_{C_j C_j} = N^{2\alpha} S_{R_j R_j}$ is unavailable and inference using this limit theory for \widehat{C}_j is infeasible. Note that under the null $C_j^0 = c_j^0 e_j$. Imposing this (maintained) restriction on the form of C_j^0 implies that the null can be rewritten as $\mathbb{H}_0 : c_j = c_j^0$ and a test analogous to the scalar case can be mounted using the j th diagonal element of the (unrestricted) estimate \widehat{C} or a similar estimate obtained by imposing the maintained restriction on C_j and estimating the system as a seemingly unrelated regression (SUR). Similar constraints on inference due to the infeasibility of the tests apply in each of these cases.

3.1.2. $c_i = c_j$ for all i, j

When $c_i = c_j = c$, for all i, j , the limiting standardized form of the signal matrix $\sum_{i=1}^N x_i x_i'$ is singular due to commonality in the explosive behavior of the components of x_t . Let $R_N = \rho_N I_K$ with $\rho_N = 1 + c/N^\alpha$. Following MP, we rotate regression coordinates to address the singularity using an orthogonal random matrix $H_N = [H_{cN}, H_{\perp N}]$ where $H_{cN} = \frac{x_N}{(x_N' x_N)^{1/2}}$ and $H_{\perp N}$ is a $K \times (K - 1)$

orthogonal complement matrix such that $H_{\perp N}' H_{cN} \stackrel{a.s.}{=} 0$. Then, $H_{\perp N}' H_{\perp N} = I_{K-1}$ and $H_{\perp N}' H_{cN} \stackrel{a.s.}{=} 0$. The limit of $H_{\perp N}$ is denoted as H_{\perp} , which satisfies $H_{\perp}' H_{\perp} = I_K - X_c X_c'$ where X_c is defined in (3.6) in Theorem 3.2. Next, rotate the regressor x_t by H_N and transform to the variate $z_t = H_N' x_t = [H_{cN}' x_t, H_{\perp N}' x_t]' = [z_{1t}', z_{2t}']'$. The following result gives the required limit theory for the LS estimator \widehat{A} in this case.

Theorem 3.2. For the discrete time system (3.1)–(3.2) with $R_N = I_K + C/N^\alpha$, $C = \text{diag}(c_1, c_2, \dots, c_K)$ with $c_i = c > 0$ for $i = 1, \dots, K$, when $N \rightarrow \infty$, we have

- (i) $\frac{1}{N^{1+\alpha}} \sum_{t=1}^N z_{2t} z_{2t}' \Rightarrow M$, with $M = H_{\perp}' \left(\frac{\mu^* \mu^{*'}}{c} + \frac{1}{2c} \Omega_{xx} \right) H_{\perp}$,
- (ii) $\frac{1}{N^{(1+\alpha)/2}} \text{vec} \left(\sum_{t=1}^N u_{0t} z_{2t}' \right) \Rightarrow \{M \otimes \Omega_{00}\}^{1/2} \times N(0, I_{m(K-1)})$, where H_{\perp} is a $K \times (K - 1)$ random matrix that is an orthogonal complement to

$$X_c = (D + \widetilde{U}_x) / \left\{ (D + \widetilde{U}_x)' (D + \widetilde{U}_x) \right\}^{1/2}, \tag{3.6}$$

satisfying $H_{\perp}H'_{\perp} = I_K - X_cX'_c$ and with

$$\begin{aligned} \tilde{U}_x &\equiv \left(\int_0^{\infty} e^{-pc} \Omega_{xx} e^{-pc} dp \right)^{1/2} U_x = \Omega_{xx}^{1/2} U_x / (2c)^{1/2}, \text{ and} \\ D &= X^* + \mu^*/c, \\ \text{(iii)} \\ N^{(1+\alpha)/2} \text{vec}(\hat{A} - A) &\Rightarrow \left(H_{\perp} M^{-1/2} \otimes \Omega_{00}^{1/2} \right) \times N(0, I_{mK}) \\ &\stackrel{d}{=} MN(0, H_{\perp} M^{-1} H'_{\perp} \otimes \Omega_{00}). \end{aligned} \tag{3.7}$$

Remark 18. When u_{0t} is weakly dependent and satisfies the conditions discussed in earlier remarks, the limit result again follows MP and has the form

$$\begin{aligned} N^{(1+\alpha)/2} \text{vec}(\hat{A} - A) &\Rightarrow \left(H_{\perp} M^{-1/2} \otimes \Omega_{00}^{1/2} \right) \times N(0, I_{mK}) \\ &\stackrel{d}{=} MN(0, H_{\perp} M^{-1} H'_{\perp} \otimes \Omega_{00}), \end{aligned} \tag{3.8}$$

where Ω_{00} is given by (3.4).

Remark 19. The limit distribution of $\hat{\mu}$ is obtained as follows:

$$\begin{aligned} &\sqrt{N}(\hat{\mu} - \mu) \\ &= \sqrt{N} \left\{ \frac{1}{N} \sum_{t=1}^N u_{xt} - \frac{1}{N} \left(\frac{1}{N^{(1+\alpha)/2}} \sum_{t=1}^N u_{xt} z'_{t-1} \right) \right. \\ &\quad \times \left. \left(\frac{1}{N^{(1+\alpha)}} \sum_{t=1}^N z_{t-1} z'_{t-1} \right)^{-1} \left(\frac{1}{N^{(1+\alpha)/2}} \sum_{t=1}^N z_{t-1} \right) \right\} \\ &\quad \left\{ \frac{N}{N} - \frac{1}{N} \left(\frac{1}{N^{(1+\alpha)/2}} \sum_{t=1}^N u_{xt} z'_{t-1} \right) \left(\frac{1}{N^{(1+\alpha)}} \sum_{t=1}^N z_{t-1} z'_{t-1} \right)^{-1} \right. \\ &\quad \times \left. \left. \left(\frac{1}{N^{(1+\alpha)/2}} \sum_{t=1}^N z_{t-1} \right) \right\}^{-1} \\ &= \frac{1}{\sqrt{N}} \sum_{t=1}^N u_{xt} + o_p(1) \Rightarrow N(0, \Omega_{xx}). \end{aligned}$$

Remark 20. Define $\hat{u}_{0t} = y_t - \hat{A}x_t$ and let the estimate of the error variance matrix be

$$S_{00} = N^{-1} \sum_{t=1}^N \hat{u}_{0t} \hat{u}'_{0t} \xrightarrow{p} \Omega_{00},$$

and the estimated variance matrix of \hat{A}_j be

$$S_{A_j A_j} = \left(\sum_{t=1}^N x_{jt}^2 \right)^{-1} S_{00}.$$

Following Theorem 3.2, we have the following limit theory for the Wald statistic for testing $\mathbb{H}_0 : Q_j A_j = Q_j A^0 = q_j$

$$W_{A_j} := \{Q_j \hat{A}_j - q_j\}' (Q_j S_{A_j A_j} Q_j)^{-1} \{Q_j \hat{A}_j - q_j\} \Rightarrow \chi_g^2.$$

Remark 21. Let $\hat{u}_{xt} = x_t - \hat{R}_N x_{t-1} - \hat{\mu}$, giving the error variance matrix estimate

$$S_{xx} = N^{-1} \sum_{t=1}^N \hat{u}_{xt} \hat{u}'_{xt} \xrightarrow{p} \Omega_{xx},$$

and the corresponding estimate of the variance matrix of \hat{R}_N (in column vector form)

$$S_{RR} = \left(\sum_{t=1}^N x_{t-1} x'_{t-1} - \frac{1}{N} \left(\sum_{t=1}^N x_{t-1} \right) \left(\sum_{t=1}^N x_{t-1} \right)' \right)^{-1} \otimes S_{xx}.$$

Then the Wald statistic for testing $\mathbb{H}_0 : Q \text{vec}(R_N) = Q \text{vec}(R_N^0) = r$, where Q is a $g \times mK$ restriction matrix of full row rank $g \leq mK$, is

$$W_{R_N} := \{Q \text{vec}(\hat{R}_N - R_N^0)\}' (Q S_{RR} Q')^{-1} \{Q \text{vec}(\hat{R}_N - R_N^0)\} \Rightarrow \chi_g^2,$$

since

$$N^{(1+\alpha)/2} Q \text{vec}(\hat{R}_N - R_N^0) \Rightarrow MN(0, Q H_{\perp} M^{-1} H'_{\perp} \otimes \Omega_{xx} Q'),$$

and

$$\begin{aligned} N^{1+\alpha} Q S_{RR} Q' &= Q \left(\frac{1}{N^{1+\alpha}} \sum_{t=1}^N x_{t-1} x'_{t-1} \right. \\ &\quad \left. - \frac{1}{N^{2+\alpha}} \left(\sum_{t=1}^N x_{t-1} \right) \left(\sum_{t=1}^N x_{t-1} \right)' \right)^{-1} \otimes S_{xx} Q' \\ &= Q H_N \left(\frac{1}{N^{1+\alpha}} \sum_{t=1}^N z_{t-1} z'_{t-1} \right)^{-1} H'_N \otimes S_{xx} Q' + o_p(1) \\ &\Rightarrow Q (H_{\perp} M^{-1} H'_{\perp} \otimes \Omega_{xx}) Q'. \end{aligned}$$

Similarly, given $R_N = 1 + \frac{c}{N^{\alpha}}$, we have $S_{CC} = N^{2\alpha} S_{RR}$ and

$$\text{vec}(\hat{C} - C^0) N^{\frac{1-\alpha}{2}} \Rightarrow MN(0, H_{\perp} M^{-1} H'_{\perp} \otimes \Omega_{xx}).$$

Hence, given α , we have the following feasible Wald test,

$$W_C := \{Q \text{vec}(\hat{C} - C^0)\}' (Q S_{CC} Q')^{-1} \{Q \text{vec}(\hat{C} - C^0)\} \Rightarrow \chi_g^2.$$

Again as in the scalar model, if α is unknown (which is the usual situation in practical work), the estimated variance matrix $S_{CC} = N^{2\alpha} S_{RR}$ is unavailable and inference using this limit theory for \hat{C} is infeasible.

Importantly, for the common explosive root case when α is known, we are able to perform statistical inference concerning the full matrix coefficients R_N and C using Wald tests because the normalization factor $N^{1+\alpha}$ is common and thereby commutable with the restriction matrix Q . However, for the distinct explosive roots case, we can only perform statistical inference about individual column vectors of R_N and C , as demonstrated in Remark 17. The same phenomenon applies for tests involving the matrix A . As shown below, these features carry over to inference in the continuous time system although in this case the sampling frequency is known so there is no difficulty relating to an unknown rate parameter α .

3.2. Limit results in the continuous time framework

The above results apply to the multivariate continuous time system

$$y(t) = \beta x(t) + u_0(t), \tag{3.9}$$

$$\begin{aligned} dx(t) &= \kappa (\mu - x(t)) dt + \Omega_{xx}^{1/2} dB_x(t), \\ x(0) &= x_0 = O_p(1), \kappa < 0, \end{aligned} \tag{3.10}$$

where $u_0(t) \sim N(0, \Omega_{00})$ is a Gaussian noise. Let $B_x(t) = \Omega_{xx}^{1/2} W_x(t)$ where W_x is standard vector Brownian motion that may be correlated with W_0 . The driver process $x(t)$ then follows a

multivariate Ornstein–Uhlenbeck process with persistence matrix κ , where $\kappa = \text{diag}(\kappa_1, \kappa_2, \dots, \kappa_K)$ is a $K \times K$ diagonal matrix. We focus on the explosive case where $\kappa_i < 0$ for $i = 1, \dots, K$. As in the discrete time case, we are interested in β , an $m \times K$ matrix of coefficients which captures co-movement between $y(t)$ and $x(t)$.

The exact discrete time representation of (3.9)–(3.10) is given by (see Phillips, 1972)

$$y_{th} = \beta x_{th} + u_{0,th}, \tag{3.11}$$

$$x_{th} = a_h(\kappa) x_{(t-1)h} + g_h + u_{x,th}, \quad x_{0h} = x_0 = O_p(1), \tag{3.11}$$

where

$$a_h(\kappa) = \exp(-\kappa h),$$

$$g_h = \kappa^{-1} (I_K - e^{-\kappa h}) \kappa \mu,$$

$$u_{x,th} = \int_{(t-1)h}^{th} e^{-\kappa(th-s)} \Omega_{xx} dB_x(s) \stackrel{d}{=} N(0, \Omega_{xx}h),$$

$$u_{0,th} \stackrel{d}{=} N(0, \Omega_{00}),$$

since

$$\mathbb{E}(u_{x,th} u'_{x,th}) = \int_{(t-1)h}^{th} e^{-2\kappa(th-s)} \Omega_{xx} ds = \frac{1}{2} \kappa^{-1} (I_K - e^{-2\kappa h}) \Omega_{xx}.$$

Thus, upon restandardization by \sqrt{h} , the system becomes

$$y_{th} = \beta x_{th} + u_{0,th}, \tag{3.12}$$

$$\tilde{x}_{th} = a_h(\kappa) \tilde{x}_{(t-1)h} + \tilde{g}_h + \tilde{u}_{x,th}, \quad \tilde{x}_{0h} = h^{-1/2} x_{0h},$$

$$\tilde{u}_{x,th} \stackrel{iid}{\sim} N(0, \Omega_{xx}), \tag{3.13}$$

where $\tilde{x}_{th} = h^{-1/2} x_{th}$, $\tilde{g}_h = h^{-1/2} g_h$ and $\tilde{u}_{x,th} = h^{-1/2} u_{x,th} \stackrel{d}{=} N(0, \Omega_{xx})$. As in the univariate case, the order of the initial value $\tilde{x}_{0h} = h^{-1/2} x_{0h}$ is $O_p(h^{-1/2})$, and the order for the drift term \tilde{g}_h is $O_p(h^{1/2})$.

For the continuous time system (3.12)–(3.13), the double asymptotic theory for the LS estimator of the coefficient matrix β when κ has distinct diagonal elements (i.e., $\kappa_i \neq \kappa_j$ for $i \neq j$) is given in the following corollary.

Corollary 3.3. Let $h = N^{-\alpha}$. For the continuous time system (3.12)–(3.13) with $\kappa = \text{diag}(\kappa_1, \kappa_2, \dots, \kappa_K)$ with $\kappa_i \neq \kappa_j$ for $i \neq j$, and $\kappa_i < 0$ for $i = 1, \dots, K$, assuming that there exists $\alpha \in (0, 1)$, such that $h^{1-\alpha} T^\alpha \rightarrow 1$ as $T \rightarrow \infty$, we have

$$\begin{aligned} & \text{vec} \left\{ \frac{1}{\sqrt{h}} (\hat{\beta} - \beta) a_h' \right\} \\ & \Rightarrow \left[\left(\int_0^\infty e^{p\kappa} (D + \tilde{U}_x) (D + \tilde{U}_x)' e^{p\kappa} dp \right)^{-1} \otimes \Omega_{00} \right]^{1/2} \\ & \quad \times N(0, I_{mK}). \end{aligned} \tag{3.14}$$

Remark 22. By multiplying both sides of (3.3) by \sqrt{h} , the double asymptotic distribution (3.14) follows directly from (3.3) with $\mu^* = \kappa \mu$, $C = -\kappa$ and $D = x_0 - \mu$. To enhance readability in terms of the relationship between the systems, we provide in Table 1 the correspondence between the models, variables and parameters in the discrete and continuous time cases.

Remark 23. The LS estimator of κ is consistent since h is known. Let $S_{00} = N^{-1} \sum_{t=1}^N \hat{u}_{0,th} \hat{u}'_{0,th}$, which satisfies

$$S_{00} \xrightarrow{p} \Omega_{00},$$

and the corresponding estimate of the covariance matrix of $\hat{\beta}_j$ is

$$S_{\beta_j \beta_j} = \left(\sum_{t=1}^N x_{jth}^2 \right)^{-1} S_{00}.$$

Table 1
Correspondence between systems (3.1)–(3.2) and (3.12)–(3.13).

Discrete time	Continuous time
$y_t = Ax_t + u_{0t}$	$\tilde{y}_{th} = \beta \tilde{x}_{th} + \tilde{u}_{0,th}$
$x_t = \mu + R_N x_{t-1} + u_{xt}$	$\tilde{x}_{th} = a_h(\kappa) \tilde{x}_{(t-1)h} + \tilde{g}_h + \tilde{u}_{x,th}$
$x_0 = x_{0N} = O_p(N^{\alpha/2})$	$\tilde{x}_{0h} = h^{-1/2} x_{0h} = O_p(h^{-1/2})$
$\mu = O_p(N^{-\alpha/2})$	$\tilde{g}_h = O_p(h^{1/2})$
C	$-\kappa$
μ with $N^{\alpha/2} \mu \rightarrow \mu^*$	\tilde{g}_h with $h^{-1/2} \tilde{g}_h \rightarrow \kappa \mu$
$X^* + C^{-1} \mu^*$	$x_0 - \mu$

The Wald statistic for testing the full rank restrictions $\mathbb{H}_0 : Q_j \beta_j = Q_j \beta_j^0 = r_j$ is then

$$W_{\beta_j} = \{Q_j (\hat{\beta}_j - \beta_j^0)\}' (Q_j S_{\beta_j \beta_j} Q_j')^{-1} \{Q_j (\hat{\beta}_j - \beta_j^0)\} \Rightarrow \chi_g^2,$$

leading to feasible inference about β_j in the continuous time framework.

Remark 24. Let a_j be the j th column of $a_h(\kappa)$. The Wald statistic for testing the full rank restrictions $\mathbb{H}_0 : Q_j a_j = Q_j a_j^0 = q_j$ for given (Q_j, q_j) has the following chi-squared limit

$$W_{a_j} := \{Q_j (\hat{a}_j - a_j^0)\}' (Q_j S_{a_j a_j} Q_j')^{-1} \{Q_j (\hat{a}_j - a_j^0)\} \Rightarrow \chi_g^2,$$

where $S_{a_j a_j} = \left(\sum_{t=1}^N x_{j(t-1)h}^2 - \frac{1}{N} \left(\sum_{t=1}^N x_{j(t-1)h} \right)^2 \right)^{-1} S_{xx}$ and $S_{xx} = N^{-1} \sum_{t=1}^N \hat{u}_{x,th} \hat{u}'_{x,th}$ satisfying $h^{-1} S_{xx} \xrightarrow{p} \Omega_{xx}$ where $\hat{u}_{x,th} = x_{th} - \hat{a}_h x_{(t-1)h} - \hat{g}_h$ are regression residuals. Let κ^j denote the j th column of κ . Given the matrix exponential relation, we have the covariance matrix of $\hat{\kappa}^j$ which satisfies $h^2 S_{\hat{\kappa}^j \hat{\kappa}^j} = S_{a_j a_j} + o(h)$ and so

$$\begin{aligned} (\hat{\kappa}^j - \kappa^{j0}) e^{-\kappa_j N} & \Rightarrow MN \left(0, \left(\int_0^\infty e^{2p\kappa_j} (D_j + \tilde{U}_{jx})^2 dp \right)^{-1} \Omega_{xx} \right) \\ & \stackrel{d}{=} MN \left(0, \frac{-2\kappa_j \Omega_{xx}}{(D_j + \tilde{U}_{jx})^2} \right). \end{aligned}$$

Then the Wald statistic for testing the (full rank) restrictions $\mathbb{H}_0 : Q_j \kappa^j = Q_j \kappa^{j0} = q_j$ satisfies

$$W_{\kappa^j} := \{Q_j \hat{\kappa}^j - q_j\}' (Q_j S_{\kappa^j \kappa^j} Q_j')^{-1} \{Q_j \hat{\kappa}^j - q_j\} \Rightarrow \chi_g^2.$$

Remark 25. The LS estimates, \hat{a}_h and $\hat{\kappa}$, do not take account of the diagonal structure of a_h and κ . If the known diagonal structure is imposed, we can use either SUR estimation or restricted LS (in which only the diagonal elements of the original LS estimates are employed). The simulation section below explores the finite sample performance of these three estimates.

Now we consider the case where the localizing explosive coefficients are identical, so that $\kappa_i = \kappa$ for $i = 1, \dots, K$.

Corollary 3.4. Let $h = N^{-\alpha}$. For the continuous time system (3.12)–(3.13) with $\kappa = \text{diag}(\kappa_1, \kappa_2, \dots, \kappa_K)$ and $\kappa_i = \bar{\kappa} < 0$ for $i = 1, \dots, K$, assume that there exists $\alpha \in (0, 1)$, such that $h^{1-\alpha} T^\alpha \rightarrow 1$ as $T \rightarrow \infty$, we have

$$\begin{aligned} \text{vec} \left\{ \sqrt{N} (\hat{\beta} - \beta) \right\} & \Rightarrow \left[H_\perp \left\{ H_\perp' \left(\mu \mu' + \frac{1}{-2\bar{\kappa}} \Omega_{xx} \right) H_\perp \right\}^{-1/2} \right. \\ & \quad \left. \otimes \Omega_{00}^{1/2} \right] \times N(0, I_{mK}). \end{aligned} \tag{3.15}$$

Remark 26. Both Corollaries 3.3 and 3.4 can be extended in the same way as before to cover the case where $u_0(t)$ is weakly dependent.

Remark 27. The double asymptotic distribution (3.15) follows directly from (3.7) with $\mu^* = \kappa\mu$ and $c = -k$.

Remark 28. The Wald statistic for testing $\mathbb{H}_0 : Qvec(\beta) = Qvec(\beta^0) = q$ for full row rank (Q, q) is then

$$W_\beta := \{Q\widehat{\beta} - q\}'(QS_{\beta\beta}Q')^{-1}\{Q\widehat{\beta} - q\} \Rightarrow \chi_g^2,$$

leading to feasible inference about the matrix coefficient β in the continuous time framework. Inference about the full matrix β is possible in this case because of the common factorization convergence rate in (3.15).

Remark 29. The Wald statistics for testing full rank restrictions on a_h and κ such as $\mathbb{H}_0 : Qvec(a_h) = Qvec(a_h^0) = q$ and $\mathbb{H}_0 : Qvec(\kappa) = Qvec(\kappa^0) = q$ are defined in a similar way and have the following chi-squared limits:

$$W_{a_h} := \{Qvec(\widehat{a}_h) - q\}'(QS_{aa}Q')^{-1}\{Qvec(\widehat{a}_h) - q\} \Rightarrow \chi_g^2,$$

and

$$W_\kappa := \{Qvec(\widehat{\kappa}) - q\}'(QS_{\kappa\kappa}Q')^{-1}\{Qvec(\widehat{\kappa}) - q\} \Rightarrow \chi_g^2,$$

again leading to feasible inference on a_h and κ because of the common factorization convergence rate.

Remark 30. When x_t has a common explosive root, LS estimation by \widehat{a}_h and $\widehat{\kappa}$ produces biased estimates due to endogeneity in the regressor, as shown in Phillips and Magdalinos (2013). The bias distorts the Wald test statistics and the distortion will be demonstrated in the Monte Carlo simulation below.

4. Monte Carlo studies

This section examines the performance of the double asymptotic limit theory in simulations. We generate data from model (2.3)-(2.4) with $\kappa = -2, \sigma_{00} = \sigma_{xx} = 1, \mu = 0$, and consider three sampling intervals, $h = 1/12, 1/52, 1/252$, corresponding to monthly, weekly and daily frequencies. The initial value x_0 is set at (0, 3, 10) and time spans of $T = 4$ and $T = 10$ years are considered. We report percentiles at levels {1%, 2.5%, 10%, 90%, 97.5%, 99%} in the limit distribution (2.21) and the finite sample distribution of the coefficient based test (called the C test hereafter) $\frac{a^N}{-2\kappa\sqrt{h}}(\widehat{\beta} - \beta)$ and t_β from Remark 9. In addition, we provide comparisons of the densities of the limit distributions and finite sample distributions of the C test statistic and t_β statistic. The number of replications is set at 10,000.

Tables 2-4 report the percentiles when $x_0 = 0, 3, 10$ by using the true values of κ and μ . It can be seen that both the double asymptotic distribution and the finite sample distribution are sensitive to changes in the initial condition for the C test. The C test is recommended for empirical studies since it provides a tighter confidence interval than the t test as demonstrated in our empirical study. In all cases the new limit distribution provides a good approximation to the finite sample distribution.

Figs. 1 and 2, and 3 plot the densities of the C test statistic and t test statistic when $T = 4$. The result is similar to the case of $T = 10$, which is not reported. These plots show the limit distribution well approximates the finite sample distribution for both tests.

To examine the performance of the limit theory in the multivariate setup, we consider a bivariate model using monthly data

($h = \frac{1}{12}$) with time span $T = 20$. Data are generated from the continuous time system (3.11), with $\beta = [1, 1]', x(0) = [3, 1]',$

$$\Omega = \begin{bmatrix} 1.5 & -0.9 & -0.8 \\ -0.9 & 2 & 0.8 \\ -0.8 & 0.8 & 1 \end{bmatrix},$$

$\mu = [1, 1]', vec(\kappa) = [\kappa_1, 0, 0, \kappa_2]'$ with $\kappa_1 = -0.2$ and $\kappa_2 = -0.4$ in the first case and $\kappa_1 = \kappa_2 = -0.2$ in the second case. Therefore, $a_1 = \exp(-\kappa_1 h) = 1.0168$ and $a_2 = 1.0339$ in the first case and $a_1 = a_2 = 1.0168$ in the second case. In Table 5, we report the percentiles of the finite sample distribution with those of the limit distribution for $W_\beta, W_\kappa^{OLS}, W_\kappa^{ReOLS}$, and W_κ^{SUR} , where W indicates the Wald test statistic for the parameter of interest, $r_\beta = [1, 1]', r_{a_h} = [1.0168, 1.0339]', r_\kappa = [-0.2, -0.4]'$ in the first case, $r_{a_h} = [1.0168, 1.0168]', r_\kappa = [-0.2, -0.2]'$ in the second case. In addition, LS corresponds to the LS estimates, ReOLS to the estimates based on the diagonal elements of LS estimates, and SUR to the estimates based on the seemingly unrelated regressions.

Several conclusions can be drawn from this Monte Carlo study. First, for β , our limit distribution well approximates the finite sample distribution in both cases. Second, for κ , the limit distribution is closer to the finite sample distribution based on SUR than those based on LS or ReOLS. While in Case 1 where the explosive roots are distinct, the finite sample distributions based on LS and SUR are very close to each other, in Case 2 where there is a common explosive root, the limit distribution is much closer to the finite sample distribution based on SUR than to those based on OLS. But in neither case was there any evidence to support the use of ReOLS. The result in Table 5 suggests that it is preferable to use the limit distribution for inference about κ based on SUR.

To understand why SUR provides much better results than LS for testing hypotheses about κ in Case 2, Table 6 reports the mean and variance of the two sets of estimates of κ in both cases. While SUR produces slightly better estimates than LS in Case 1, it yields much better estimates of κ in Case 2. As shown in Phillips and Magdalinos (2013), due to the endogeneity problem in the VAR models when there is a common explosive root, the LS estimate of the common explosive autoregression parameter is biased downward, suggesting that the estimate of κ is biased upward. Naturally this bias distorts the asymptotic approximation of the Wald statistic.

5. Empirical illustration for the US real estate market

This section illustrates use of the limit theory in an empirical study of the relationship between the US nationwide real estate market and metropolitan real estate markets, respectively, between January 2000 and April 2006. We apply the limit theory for univariate co-moving system (2.1)-(2.2) to real estate data using the S&P/Case-Shiller home price composite 20-city index and metropolitan area indices. The S&P/Case-Shiller home price indices are the leading measures of the US residential real estate prices, tracking changes in the value of residential real estate nationwide. Monthly data for the composite 20-city index and 20 metropolitan area indices between February 2000 and August 2014 were downloaded from the St. Louis Fed.³

Similar to the capital asset pricing model CAPM, we use the composite 20 index to measure the overall market movements. A multi-equation continuous time system (2.1)-(2.2) is estimated with x_t as the logarithmic composite 20 index (called the countrywide index in the present paper) and each y_t being one of the logarithmic metropolitan area indices that is found to be explosive. The coefficient β then measures the co-movement of each metropolitan area index with the countrywide index. With monthly data, the sampling interval is set to $h = 1/12$. The initial

³ <http://research.stlouisfed.org/fred2/release?rid=199>.




Fig. 1. Density comparison between the C test and t test for both finite sample and limit distributions, when the initial value is $x_0 = 0$.




Fig. 2. Density comparison between the C test and t test for both finite sample and limit distributions, when the initial value is $x_0 = 3$.

Table 2
Comparison of the finite sample and double asymptotic distributions of $\hat{\beta}$, when the initial value is $x_0 = 0$.

Time Span		T = 4						T = 10					
Frequency	C test	1%	2.50%	10%	90%	97.50%	99%	1%	2.50%	10%	90%	97.50%	99%
Daily (h = 1/252)	New	-32.233	-13.222	-3.027	3.063	12.063	29.452	-32.233	-13.222	-3.027	3.063	12.063	29.452
	Finite Sample	-36.346	-13.501	-3.271	2.870	11.647	30.428	-28.528	-12.100	-2.889	3.166	13.304	32.007
Weekly (h = 1/52)	New	-32.233	-13.222	-3.027	3.063	12.063	29.452	-32.233	-13.222	-3.027	3.063	12.063	29.452
	Finite Sample	-31.973	-11.329	-3.006	2.973	11.752	27.031	-32.775	-12.919	-3.145	3.018	12.735	35.756
Monthly (h = 1/12)	New	-32.233	-13.222	-3.027	3.063	12.063	29.452	-32.233	-13.222	-3.027	3.063	12.063	29.452
	Finite Sample	-34.772	-12.327	-2.889	2.793	12.284	27.255	-31.209	-11.698	-2.922	2.842	11.947	32.185
t test													
Daily (h = 1/252)	New	-2.326	-1.960	-1.282	1.282	1.960	2.326	-2.326	-1.960	-1.282	1.282	1.960	2.326
	Finite Sample	-2.305	-1.950	-1.271	1.253	1.915	2.256	-2.380	-1.976	-1.277	1.301	2.015	2.359
Weekly (h = 1/52)	New	-2.326	-1.960	-1.282	1.282	1.960	2.326	-2.326	-1.960	-1.282	1.282	1.960	2.326
	Finite Sample	-2.334	-2.004	-1.292	1.318	1.956	2.340	-2.375	-2.000	-1.286	1.248	1.936	2.285
Monthly (h = 1/12)	New	-2.326	-1.960	-1.282	1.282	1.960	2.326	-2.326	-1.960	-1.282	1.282	1.960	2.326
	Finite Sample	-2.452	-2.072	-1.326	1.331	2.057	2.463	-2.434	-1.977	-1.301	1.306	1.969	2.301

Table 3
Comparison of the finite sample and double asymptotic distributions of $\hat{\beta}$, when the initial value is $x_0 = 3$.

Time Span		T = 4						T = 10					
Frequency	C test	1%	2.50%	10%	90%	97.50%	99%	1%	2.50%	10%	90%	97.50%	99%
Daily (h = 1/252)	New	-0.427	-0.348	-0.220	0.219	0.347	0.426	-0.427	-0.348	-0.220	0.219	0.347	0.426
	Finite Sample	-0.416	-0.338	-0.215	0.215	0.341	0.412	-0.426	-0.350	-0.220	0.218	0.346	0.418
Weekly (h = 1/52)	New	-0.427	-0.348	-0.220	0.219	0.347	0.426	-0.427	-0.348	-0.220	0.219	0.347	0.426
	Finite Sample	-0.413	-0.338	-0.215	0.215	0.337	0.414	-0.411	-0.341	-0.218	0.212	0.336	0.413
Monthly (h = 1/12)	New	-0.427	-0.348	-0.220	0.219	0.347	0.426	-0.427	-0.348	-0.220	0.219	0.347	0.426
	Finite Sample	-0.385	-0.324	-0.204	0.205	0.321	0.386	-0.391	-0.318	-0.200	0.207	0.322	0.391
t test													
Daily (h = 1/252)	New	-2.326	-1.960	-1.282	1.282	1.960	2.326	-2.326	-1.960	-1.282	1.282	1.960	2.326
	Finite Sample	-2.275	-1.924	-1.271	1.249	1.947	2.283	-2.369	-1.989	-1.290	1.285	1.993	2.358
Weekly (h = 1/52)	New	-2.326	-1.960	-1.282	1.282	1.960	2.326	-2.326	-1.960	-1.282	1.282	1.960	2.326
	Finite Sample	-2.331	-1.976	-1.309	1.300	1.985	2.340	-2.352	-1.965	-1.285	1.249	1.950	2.320
Monthly (h = 1/12)	New	-2.326	-1.960	-1.282	1.282	1.960	2.326	-2.326	-1.960	-1.282	1.282	1.960	2.326
	Finite Sample	-2.439	-2.072	-1.325	1.327	2.050	2.469	-2.397	-1.986	-1.293	1.311	1.963	2.364

Table 4
Comparison of the finite sample and double asymptotic distributions of $\hat{\beta}$, when the initial value is $x_0 = 10$.

Time Span		T = 4						T = 10					
Frequency	C test	1%	2.50%	10%	90%	97.50%	99%	1%	2.50%	10%	90%	97.50%	99%
Daily (h = 1/252)	New	-0.118	-0.099	-0.064	0.064	0.099	0.117	-0.118	-0.099	-0.064	0.064	0.099	0.117
	Finite Sample	-0.114	-0.096	-0.063	0.063	0.098	0.115	-0.119	-0.100	-0.064	0.064	0.099	0.117
Weekly (h = 1/52)	New	-0.118	-0.099	-0.064	0.064	0.099	0.117	-0.118	-0.099	-0.064	0.064	0.099	0.117
	Finite Sample	-0.114	-0.097	-0.064	0.063	0.097	0.116	-0.116	-0.096	-0.063	0.061	0.094	0.115
Monthly (h = 1/12)	New	-0.118	-0.099	-0.064	0.064	0.099	0.117	-0.118	-0.099	-0.064	0.064	0.099	0.117
	Finite Sample	-0.109	-0.092	-0.059	0.060	0.092	0.108	-0.110	-0.091	-0.059	0.060	0.090	0.110
t test													
Daily (h = 1/252)	New	-2.326	-1.960	-1.282	1.282	1.960	2.326	-2.326	-1.960	-1.282	1.282	1.960	2.326
	Finite Sample	-2.274	-1.924	-1.271	1.249	1.947	2.283	-2.369	-1.989	-1.290	1.285	1.993	2.358
Weekly (h = 1/52)	New	-2.326	-1.960	-1.282	1.282	1.960	2.326	-2.326	-1.960	-1.282	1.282	1.960	2.326
	Finite Sample	-2.331	-1.976	-1.309	1.300	1.985	2.340	-2.352	-1.965	-1.285	1.249	1.950	2.320
Monthly (h = 1/12)	New	-2.326	-1.960	-1.282	1.282	1.960	2.326	-2.326	-1.960	-1.282	1.282	1.960	2.326
	Finite Sample	-2.439	-2.072	-1.325	1.327	2.050	2.469	-2.397	-1.986	-1.293	1.311	1.963	2.364

Table 5
Comparison of the finite sample and double asymptotic distributions of the Wald tests related to $\hat{\beta}_{2 \times 1}$ and $\hat{\kappa}_{2 \times 1}$.

Wald test	Percentile	Case 1: $\kappa_1 = -0.2, \kappa_2 = -0.4$						Case 2: $\kappa_1 = -0.2, \kappa_2 = -0.2$					
		1%	2.50%	10%	90%	97.50%	99%	1%	2.50%	10%	90%	97.50%	99%
Asymptotic	χ_g^2	0.020	0.051	0.211	4.605	7.378	9.210	0.020	0.051	0.211	4.605	7.378	9.210
W_{β}	Finite	0.023	0.057	0.206	4.626	7.373	9.354	0.021	0.055	0.226	4.831	7.871	9.760
W_{β}^{OLS}	Finite	0.020	0.048	0.209	5.264	9.215	12.752	7.645	8.953	11.391	20.142	24.165	26.482
W_{κ}^{ReOLS}	Finite	0.000	0.000	0.000	0.044	1.466	7.980	5.035	6.104	8.082	17.906	22.410	24.899
W_{κ}^{SUR}	Finite	0.020	0.050	0.201	4.589	7.509	9.688	0.021	0.050	0.225	5.087	8.215	10.347




Fig. 3. Density comparison between the C test and t test for both finite sample and limit distributions, when the initial value is $x_0 = 10$.

Table 6
Finite sample comparison of $\hat{\beta}_{2 \times 1}$ and $\hat{\kappa}_{2 \times 1}$ for the OLS and SUR estimates.

Method	Case 1: $\kappa_1 = -0.2, \kappa_2 = -0.4$					Case 2: $\kappa_1 = -0.2, \kappa_2 = -0.2$				
	TRUE	Mean	VAR	Mean	VAR	TRUE	Mean	VAR	Mean	VAR
κ_1	-0.200	-0.187	4.60E-03	-0.197	7.16E-04	-0.200	0.042	9.03E-02	-0.197	7.50E-04
κ_2	-0.400	-0.400	1.38E-05	-0.400	2.17E-07	-0.200	0.200	1.05E-01	-0.195	1.12E-03

value in each equation of the system is set to the composite 20 index in January 2000, i.e., $x_0 = \log(100.59) = 4.61$.

We focus on the sample period between January 2000 and April 2006 (in this case $T = 6.25$). The choice of the sample period is guided by recent work in Phillips and Yu, (2011) that documents the presence of the explosive behavior in the US real estate market over much of this period. Before estimating the main model (2.1), we first explore the presence of explosive behavior in the market index (x_{it}) and in the individual area indices (y_{it}) by estimating κ and κ_y and obtaining the standard errors, the t-statistics, and the 99% and 90% confidence intervals based on the C test and the t test using the double asymptotic theory developed in WY. Results are reported in Table 7. For the countrywide index, the LS estimate of κ is -0.0467 , with the estimated standard error of 0.0002 and the t statistic of -2.3538 . Its 90% confidence interval is $[-0.0486, -0.0449]$ based on the C test and $[-0.0793, -0.0141]$ based on the t test. All the results confirm the explosive behavior in x_{it} over this period. For 11 out of 20 individual area indices, we find explosive behavior based on both the 90% and the 99% confidence intervals by the C test. Moreover, upon comparison of the confidence intervals reported in Table 7, it is apparent that the C test produces tighter confidence intervals than the t test.

For the 11 areas that exhibit explosive behavior, we further study possible co-movement with the market index. We report the

LS estimate of β , the estimated standard error, the t-statistic, and the 99% and 90% confidence intervals based on the C test and the t test in Table 8. Allowing for possible weak dependence in u_{0t} and following Remark 7 we estimate the variance and long-run variances of u_{0t} by

$$\hat{\sigma}_{00}^2 = \frac{1}{N} \sum_{h=1}^N \hat{u}_{0t} \hat{u}_{0t-h}^2, \quad \hat{\omega}_{00}^2 = \hat{\sigma}_{00}^2 + 2\hat{\lambda}_{00}^2,$$

$$\hat{\lambda}_{00}^2 = \frac{1}{N} \sum_{h=1}^{M_n} \left(1 - \frac{h}{M_n + 1}\right) \sum_{t=h+1}^N \hat{u}_{0t} \hat{u}_{0t-h}$$

with truncation lag $M_n = N^{1/3}$. Evidently, the 90% confidence intervals are quite tight and comfortably reject the null hypothesis $H_0 : \beta = 0$ in all cases. The confidence intervals can also be used to assess whether $\beta = 1$ versus $\beta < 1$ or $\beta > 1$. If $\beta > 1$ (respectively, $\beta < 1$) the index of the associated metropolitan area moves faster (slower) than the countrywide index, giving a useful perspective on the relationship of different metropolitan area indices to the national index. The results show that LA, Miami, DC, and NY have more “aggressive” real estate markets in the US than the whole market, in the sense that the index for these cities moves more than the countrywide index.

Table 7

The estimated persistence parameter in y_{it} and x_{it} , the confidence intervals for persistence parameter in the US logarithmic real estate market.

Place	κ or κ_y	Std Err	t-stat	C test 99% CI		C test 90% CI		t test 99% CI		t test 90% CI	
Market	-0.0467	1.9843E-02	-2.3538	-0.0496	-0.0438	-0.0486	-0.0449	-0.0978	0.0044	-0.0793	-0.0141
SF	0.0669	6.8910E-02	0.9705	0.0753	0.0584	0.0723	0.0615	-0.1106	0.2444	-0.0465	0.1802
LA	-0.0871	2.5867E-02	-3.3658	-0.0913	-0.0829	-0.0897	-0.0844	-0.1537	-0.0204	-0.1296	-0.0445
LasVegas	-0.1072	5.9086E-02	-1.8146	-0.1118	-0.1027	-0.1101	-0.1043	-0.2594	0.0450	-0.2044	-0.0100
Miami	-0.1605	1.8272E-02	-8.7839	-0.1654	-0.1556	-0.1636	-0.1574	-0.2076	-0.1134	-0.1906	-0.1304
Phoenix	-0.3464	4.9243E-02	-7.0355	-0.3497	-0.3431	-0.3486	-0.3443	-0.4733	-0.2196	-0.4274	-0.2655
SanDiego	0.0526	3.3438E-02	1.5729	0.0587	0.0465	0.0565	0.0487	-0.0335	0.1387	-0.0024	0.1076
Denver	0.4462	5.7549E-02	7.7529	1.0475	-0.1569	0.8304	0.0617	0.2979	0.5944	0.3515	0.5408
DC	-0.0536	2.3675E-02	-2.2647	-0.0568	-0.0504	-0.0556	-0.0516	-0.1146	0.0074	-0.0926	-0.0147
Chicago	-0.0199	2.8680E-02	-0.6929	-0.0213	-0.0184	-0.0208	-0.0189	-0.0937	0.0540	-0.0670	0.0273
Boston	0.2315	3.2179E-02	7.1950	0.3131	0.1497	0.2837	0.1794	0.1486	0.3144	0.1786	0.2845
Charlotte	-0.1667	1.2529E-01	-1.3307	-0.1716	-0.1618	-0.1698	-0.1636	-0.4894	0.1560	-0.3728	0.0394
Portland	-0.3691	3.2029E-02	-115238	-0.3721	-0.3660	-0.3710	-0.3671	-0.4516	-0.2866	-0.4218	-0.3164
Dallas	0.2945	1.1385E-01	2.5866	0.4483	0.1402	0.3928	0.1962	0.0012	0.5878	0.1072	0.4818
Detroit	0.2508	7.3621E-02	3.4072	0.3506	0.1508	0.3146	0.1871	0.0612	0.4405	0.1297	0.3719
Atlanta	0.0453	4.3169E-02	1.0486	0.0502	0.0403	0.0484	0.0421	-0.0659	0.1565	-0.0257	0.1163
Seattle	-0.2937	2.8503E-02	-103035	-0.2976	-0.2898	-0.2962	-0.2912	-0.3671	-0.2203	-0.3406	-0.2468
Minneapolis	0.1418	2.6267E-02	5.4002	0.1704	0.1132	0.1601	0.1236	0.0742	0.2095	0.0986	0.1850
Tampa	-0.2111	2.9201E-02	-7.2302	-0.2158	-0.2064	-0.2141	-0.2081	-0.2863	-0.1359	-0.2592	-0.1631
Cleveland	0.1000	6.5902E-02	1.5175	0.1155	0.0845	0.1099	0.0901	-0.0697	0.2698	-0.0084	0.2084
NY	-0.0005	1.9307E-02	-0.0242	-0.0005	-0.0004	-0.0005	-0.0004	-0.0502	0.0493	-0.0322	0.0313

Table 8

The estimated β coefficients and the confidence intervals for β in the US logarithmic real estate market.

City	β	S.E(β)	t(β)	C test 99% CI		C test 90% CI		t test 99% CI		t test 90% CI	
LA	1.0223	0.0052	4.3232	1.0046	1.0401	1.0110	1.0337	1.0090	1.0357	1.0138	1.0308
LasVegas	1.0009	0.0053	0.1796	0.9828	1.0191	0.9894	1.0125	0.9873	1.0146	0.9923	1.0096
Miami	1.0163	0.0045	3.6168	1.0008	1.0318	1.0064	1.0261	1.0047	1.0279	1.0089	1.0237
Phoenix	0.9832	0.0042	-4.0050	0.9688	0.9976	0.9740	0.9924	0.9724	0.9940	0.9763	0.9901
DC	1.0190	0.0034	5.6085	1.0074	1.0307	1.0116	1.0264	1.0103	1.0277	1.0134	1.0246
Chicago	0.9806	0.0033	-5.8713	0.9693	0.9920	0.9734	0.9879	0.9722	0.9891	0.9752	0.9861
Charlotte	0.9456	0.0075	-7.2537	0.9199	0.9714	0.9292	0.9621	0.9263	0.9649	0.9333	0.9580
Portland	0.9670	0.0035	-9.5031	0.9551	0.9790	0.9594	0.9746	0.9581	0.9760	0.9613	0.9727
Seattle	0.9713	0.0036	-7.9975	0.9590	0.9837	0.9635	0.9792	0.9621	0.9806	0.9654	0.9772
Tampa	1.0000	0.0021	0.0089	0.9927	1.0074	0.9953	1.0047	0.9945	1.0055	0.9965	1.0035
NY	1.0083	0.0013	6.5547	1.0039	1.0126	1.0055	1.0111	1.0050	1.0115	1.0062	1.0104

6. Conclusion

This paper studies co-moving systems with explosive regressors in a continuous time framework. The exact discretized model corresponds to a modified version of the discrete time model of MP but allows for larger initial condition effects and an asymptotically negligible intercept. The limit theory is developed for this modified model, enabling us to obtain double asymptotic limit theory for a continuous time system in which the span $T \rightarrow \infty$ and the sampling interval $h \rightarrow 0$. The extensions have some important implications for practical work. First, the limit distribution depends explicitly on the initial condition. This dependence mimics a corresponding property in the finite sample distribution and thereby improves the quality of the double asymptotic limit theory as a finite sample approximation. Second, the localized coefficient c , whose counterpart in continuous time is $-\kappa$, is consistently estimable in continuous time using the LS estimator, facilitating a coefficient based test for mildly explosive behavior. Finally, pivotal inference is facilitated in the continuous time case because the sampling interval is known whereas in discrete time system the corresponding localizing rate parameter is unknown.

The double asymptotic limit theory is developed for univariate and multivariate discrete time systems that relate to systems formulated in continuous time. Simulations suggest that for the coefficient based test and the t test statistics, these asymptotics well approximate the finite sample distributions. An empirical illustration with US real estate prices at national and various metropolitan areas shows how the methods assist in identifying regions where real estate markets are more aggressive than others.

Appendix A

A.1. Proof of Theorem 2.1

Proof. The arguments here and in much of what follows closely mirror those of MP in the mildly explosive case. We therefore provide only the main new details here. The limit theory of $\sum_{t=1}^N x_t^2$ and $\sum_{t=1}^N x_t u_{0t}$ is obtained using split sample arguments replacing summations in $\sum_{t=1}^N$ by $(\sum_{t=1}^{m_N} + \sum_{t=m_N+1}^N)$ where m_N is such that $\frac{m_N}{N^\alpha} + \frac{N}{m_N} \rightarrow \infty$ so that with $c > 0$ and $\alpha \in (0, 1)$ we have

$$R_N^{-m_N} \sim \left(1 + \frac{c}{N^\alpha}\right)^{-m_N} \rightarrow 0, \quad \frac{N^\alpha}{R_N^{N-m_N}} \rightarrow 0. \tag{A.1}$$

(i) Start by writing x_t in (2.9) as follows:

$$x_t = R_N^t x_0 + \sum_{j=1}^t R_N^{t-j} u_{xj} + \frac{1 - R_N^t}{1 - R_N} \mu, \tag{A.2}$$

so the standardized numerator can be decomposed as

$$\begin{aligned} & (R_N^N N^\alpha)^{-1} \sum_{t=1}^N x_t u_{0t} \\ &= (R_N^N N^\alpha)^{-1} \sum_{t=1}^N R_N^t u_{0t} \left(x_0 - \frac{\mu}{1 - R_N}\right) \\ &+ (R_N^N N^\alpha)^{-1} \sum_{t=1}^N u_{0t} \frac{\mu}{1 - R_N} \end{aligned}$$

$$+ \frac{R_N^{-N}}{\sqrt{N^\alpha}} \left(\sum_{t=1}^{m_N} + \sum_{t=m_N+1}^N \right) u_{0t} \left(\frac{1}{\sqrt{N^\alpha}} \sum_{j=1}^t R_N^{t-j} u_{xj} \right). \quad (\text{A.3})$$

For the first term on the right hand side of (A.3), since $\tilde{x}_0 = x_{0N} N^{-\alpha/2} \Rightarrow X^*$ and $\tilde{\mu} = N^{\alpha/2} \mu \Rightarrow \mu^*$, we have

$$D_N = N^{-\alpha/2} x_0 - \frac{N^{-\alpha/2} \mu}{1 - R_N} = N^{-\alpha/2} x_0 - \frac{N^{\alpha/2} \mu}{-c + o(1)} \\ \Rightarrow X^* + \frac{\mu^*}{c} = D. \quad (\text{A.4})$$

Setting $D_N = N^{-\alpha/2} \left(x_0 - \frac{\mu}{1 - R_N} \right)$, we then have

$$\begin{aligned} & (R_N^N N^\alpha)^{-1} \sum_{t=1}^N R_N^t u_{0t} \left(x_0 - \frac{\mu}{1 - R_N} \right) \\ &= N^{-\alpha/2} \sum_{t=1}^N R_N^{-(N-t)} u_{0t} \left\{ N^{-\alpha/2} \left(x_0 - \frac{\mu}{1 - R_N} \right) \right\} \\ &= D_N N^{-\alpha/2} \sum_{t=1}^N R_N^{-(N-t)} u_{0t} \\ &= D N^{-\alpha/2} \sum_{t=m_N+1}^N R_N^{-(N-t)} u_{0t} + o_p(1), \text{ with } D := X^* + \frac{\mu^*}{c}, \end{aligned} \quad (\text{A.5})$$

where we assume the probability space is expanded in such a way so that the weak convergence \Rightarrow can be replaced by \xrightarrow{p} . Also note that

$$\begin{aligned} \mathbb{E} \left\{ N^{-\alpha/2} \sum_{t=1}^{m_N} R_N^{-(N-t)} u_{0t} \right\}^2 &= \sigma_{00}^2 N^{-\alpha} \sum_{t=1}^{m_N} R_N^{-2(N-t)} \\ &= \sigma_{00}^2 N^{-\alpha} R_N^{-2(N+1)} \frac{1 - R_N^{2m_N}}{1 - R_N^2} \\ &= \sigma_{00}^2 \frac{R_N^{-2(N+1)} - R_N^{-2(N+1)+2m_N}}{-2c} \\ &= o(1), \end{aligned}$$

so that

$$N^{-\alpha/2} \sum_{t=1}^{m_N} R_N^{-(N-t)} u_{0t} = o_p(1), \quad (\text{A.6})$$

and then

$$(R_N^N N^\alpha)^{-1} \sum_{t=1}^{m_N} R_N^t u_{0t} \left(x_0 - \frac{\mu}{1 - R_N} \right) = o_p(1).$$

Hence, for the first term of (A.3) we have by virtue of the martingale central limit theorem (MCLT), as in PM,

$$\begin{aligned} & (R_N^N N^\alpha)^{-1} \sum_{t=1}^N R_N^t u_{0t} \left(x_0 - \frac{\mu}{1 - R_N} \right) \\ &= D_N (R_N^N N^{\alpha/2})^{-1} \sum_{t=1}^N R_N^t u_{0t} \\ &= D \frac{1}{\sqrt{N^\alpha}} \sum_{t=m_N+1}^N R_N^{-(N-t)} u_{0t} + o_p(1) \\ &= D \left(\frac{1}{\sqrt{N^\alpha}} \sum_{k=0}^{N-m_N-1} R_N^{-k} u_{0N-k} \right) \\ &= D \left(\frac{1}{\sqrt{N^\alpha}} \sum_{k=0}^{N-m_N-1} R_N^{-k} u'_{0k} \right) \Rightarrow \frac{D \sigma_{00}}{(2c)^{1/2}} U_0, \end{aligned}$$

where $u'_{0k} := u_{0N-k} \stackrel{iid}{\sim} (0, \sigma_{00}^2)$ and $U_0 = N(0, 1)$ since $\frac{1}{N^\alpha} \sum_{k=0}^{N-m_N-1} R_N^{-2k} = \frac{1}{N^\alpha} \frac{1 - R_N^{-2N}}{1 - R_N^{-2}} \sim \frac{1}{N^\alpha} \frac{1}{R_N^{-2}} \rightarrow \frac{1}{2c}$.

For the second term on the right hand side of (A.3), noting that $R_N^{-N} \sqrt{N} = (1 + \frac{c}{N^\alpha})^{-N} \sqrt{N} = o\left(e^{-c \frac{N}{N^\alpha}} \sqrt{N}\right) = o(1)$ for all $\alpha \in (0, 1)$ we obtain

$$\begin{aligned} & (R_N^N N^\alpha)^{-1} \sum_{t=1}^N u_{0t} \frac{\mu}{1 - R_N} \sim R_N^{-N} \sqrt{N} \frac{\mu}{-c} \frac{1}{\sqrt{N}} \sum_{t=1}^N u_{0t} \\ &= O\left(e^{-c \frac{N}{N^\alpha}} \sqrt{\frac{N}{N^\alpha}}\right) O_p(1) = o_p(1). \end{aligned} \quad (\text{A.7})$$

The third term on the right hand side of (A.3) is

$$\begin{aligned} & \frac{R_N^{-N}}{\sqrt{N^\alpha}} \left(\sum_{t=1}^{m_N} + \sum_{t=m_N+1}^N \right) u_{0t} \left(\frac{1}{\sqrt{N^\alpha}} \sum_{j=1}^t R_N^{t-j} u_{xj} \right) \\ &= \frac{1}{\sqrt{N^\alpha}} \sum_{t=1}^{m_N} R_N^{-(N-t)} u_{0t} \left(\frac{1}{\sqrt{N^\alpha}} \sum_{j=1}^t R_N^{-j} u_{xj} \right) \\ &+ \frac{1}{\sqrt{N^\alpha}} \sum_{t=m_N+1}^N R_N^{-(N-t)} u_{0t} \left(\frac{1}{\sqrt{N^\alpha}} \sum_{j=1}^t R_N^{-j} u_{xt} \right) \\ &= \frac{1}{\sqrt{N^\alpha}} \sum_{t=m_N+1}^N R_N^{-(N-t)} u_{0t} \left(\frac{1}{\sqrt{N^\alpha}} \sum_{j=1}^{m_N} R_N^{-j} u_{xj} \right) + o_p(1), \end{aligned}$$

where we use the fact that $N^{-\alpha/2} \sum_{t=1}^{m_N} R_N^{-(N-t)} u_{0t} = o_p(1)$ from (A.6). We now use a joint MCLT for the components

$$\begin{aligned} (U_{0N}, U_{xN}) &= \left(\frac{1}{\sqrt{N^\alpha}} \sum_{t=m_N+1}^N R_N^{-(N-t)} u_{0t}, \frac{1}{\sqrt{N^\alpha}} \sum_{j=1}^{m_N} R_N^{-j} u_{xj} \right) \\ &= \left(\frac{1}{\sqrt{N^\alpha}} \sum_{t=1}^N R_N^{-(N-t)} u_{0t}, \frac{1}{\sqrt{N^\alpha}} \sum_{j=1}^{m_N} R_N^{-j} u_{xj} \right) + o_p(1) \\ &\Rightarrow \left(\frac{\sigma_{00}}{(2c)^{1/2}} U_0, \frac{\sigma_{xx}}{(2c)^{1/2}} U_x \right) \text{ with } (U'_0, U'_x)' \\ &\sim N(0, I_2), \end{aligned}$$

just as in PM and MP, using the fact that the limit variates (U_0, U_x) are independent because

$$\begin{aligned} & \mathbb{E} \left\{ \left(\frac{1}{\sqrt{N^\alpha}} \sum_{t=1}^N R_N^{-(N-t)} u_{0t} \right) \left(\frac{1}{\sqrt{N^\alpha}} \sum_{j=1}^{m_N} R_N^{-j} u_{xj} \right) \right\} \\ &= \frac{N^{1-\alpha}}{R_N^N} \sigma_{0x} \rightarrow 0. \end{aligned}$$

Hence

$$\begin{aligned} & \frac{R_N^{-N}}{\sqrt{N^\alpha}} \left(\sum_{t=1}^{m_N} + \sum_{t=m_N+1}^N \right) u_{0t} \left(\frac{1}{\sqrt{N^\alpha}} \sum_{j=1}^t R_N^{t-j} u_{xj} \right) \\ &= \frac{1}{\sqrt{N^\alpha}} \sum_{t=m_N+1}^N R_N^{-(N-t)} u_{0t} \left(\frac{1}{\sqrt{N^\alpha}} \sum_{j=1}^{m_N} R_N^{-j} u_{xj} \right) + o_p(1) \\ &\Rightarrow \left(\frac{\sigma_{00}}{(2c)^{1/2}} U_0 \right) \left(\frac{\sigma_{xx}}{(2c)^{1/2}} U_x \right) = \frac{\sigma_{00} \sigma_{xx}}{2c} U_0 U_x. \end{aligned}$$

Combining the above results and using (A.4) we obtain

$$(R_N^N N^\alpha)^{-1} \sum_{t=1}^N x_t u_{0t}$$

$$\begin{aligned}
 &= (R_N^N N^\alpha)^{-1} \sum_{t=1}^N R_N^t u_{0t} \left(x_0 - \frac{\mu}{1 - R_N} \right) \\
 &\quad + (R_N^N N^\alpha)^{-1} \sum_{t=1}^N u_{0t} \frac{\mu}{1 - R_N} \\
 &\quad + \frac{R_N^{-N}}{\sqrt{N^\alpha}} \sum_{t=m_N+1}^N u_{0t} \left(\frac{1}{\sqrt{N^\alpha}} \sum_{j=1}^t R_N^{t-j} u_{xj} \right) + o_p(1) \\
 &= (R_N^N N^\alpha)^{-1} \sum_{t=1}^N R_N^t u_{0t} \left(x_0 - \frac{\mu}{1 - R_N} \right) \\
 &\quad + \frac{R_N^{-N}}{\sqrt{N^\alpha}} \sum_{t=m_N+1}^N u_{0t} \left(\frac{1}{\sqrt{N^\alpha}} \sum_{j=1}^t R_N^{t-j} u_{xj} \right) + o_p(1) \\
 &\Rightarrow \frac{D\sigma_{00}}{(2c)^{1/2}} U_0 + \frac{\sigma_{00}\sigma_{xx}}{2c} U_0 U_x \\
 &= \frac{\sigma_{00}}{(2c)^{1/2}} U_0 \left(D + \frac{\sigma_{xx}}{(2c)^{1/2}} U_x \right), \tag{A.8}
 \end{aligned}$$

giving the limit of the numerator.

(ii) From the identity

$$x_t^2 = R_N^2 x_{t-1}^2 + \mu^2 + u_{xt}^2 + 2R_N \mu x_{t-1} + 2R_N x_{t-1} u_{xt} + 2\mu u_{xt},$$

we have

$$\begin{aligned}
 (R_N^2 - 1) \sum_{t=1}^N x_t^2 &= R_N^2 x_N^2 - R_N^2 x_0^2 - 2R_N \mu \sum_{t=1}^N x_{t-1} \\
 &\quad - 2R_N \sum_{t=1}^N x_{t-1} u_{xt} - N\mu^2 - \sum_{t=1}^N u_{xt}^2 - 2\mu \sum_{t=1}^N u_{xt}. \tag{A.9}
 \end{aligned}$$

We show in the following that each of the following standardized terms

$$\begin{aligned}
 &\frac{R_N^2 x_0^2}{R_N^2 N^\alpha}, \frac{N\mu^2}{R_N^2 N^\alpha}, \frac{2\mu \sum_{t=1}^N u_{xt}}{R_N^2 N^\alpha}, \frac{\sum_{t=1}^N u_{xt}^2}{R_N^2 N^\alpha}, \\
 &\frac{R_N \mu \sum_{t=1}^N x_{t-1}}{R_N^2 N^\alpha}, \frac{R_N \sum_{t=1}^N x_{t-1} u_{xt}}{R_N^2 N^\alpha}
 \end{aligned}$$

are asymptotically negligible. In particular, since the standardized initial condition and drift satisfy $\tilde{x}_0 = x_{0N} N^{-\alpha/2} \Rightarrow X^*$ and $\tilde{\mu} = N^{\alpha/2} \mu \Rightarrow \mu^*$ we find that

$$\frac{R_N^2 x_0^2}{R_N^2 N^\alpha} = O_p \left(\left(\frac{x_0}{N^{\alpha/2}} \right)^2 \frac{1}{R_N^2} \right) = o_p(1),$$

$$\frac{N\mu^2}{R_N^2 N^\alpha} = O_p \left(\frac{N^{1-2\alpha}}{R_N^2} \right) = o_p(1),$$

$$\begin{aligned}
 \frac{2\mu \sum_{t=1}^N u_{xt}}{R_N^2 N^\alpha} &= \left(\frac{1}{R_N^2} \right) \left(\frac{2\mu\sqrt{N}}{N^\alpha} \right) \left(\frac{1}{\sqrt{N}} \sum_{t=1}^N u_{xt} \right) \\
 &= O_p \left(\frac{1}{R_N^2} \right) \times O_p \left(N^{\frac{1}{2}-\frac{3}{2}\alpha} \right) \times O_p(1) = o_p(1),
 \end{aligned}$$

$$\frac{\sum_{t=1}^N u_{xt}^2}{R_N^2 N^\alpha} = \frac{N}{R_N^2 N^\alpha} \frac{1}{N} \sum_{t=1}^N u_{xt}^2 = O_p \left(\frac{N^{1-\alpha}}{R_N^2} \right) \times O_p(1) = o_p(1),$$

$$\frac{R_N \sum_{t=1}^N x_{t-1} u_{xt}}{R_N^2 N^\alpha} = O_p \left(\frac{\sum_{t=1}^N x_{t-1} u_{xt}}{R_N^2 N^\alpha} \right) \times O_p \left(\frac{1}{R_N^2} \right) = o_p(1),$$

since $(R_N^N N^\alpha)^{-1} \sum_{t=1}^N x_{t-1} u_{xt} = O_p(1)$ just as in the analysis of $(R_N^N N^\alpha)^{-1} \sum_{t=1}^N x_t u_{0t}$ in part (i); and finally

$$\begin{aligned}
 \frac{R_N \mu \sum_{t=1}^N x_t}{R_N^2 N^\alpha} &= \frac{\mu}{N^{\alpha/2} R_N^2} \sum_{t=1}^N \frac{x_t}{N^{\alpha/2} R_N^2} \frac{R_N^t}{R_N^2} \\
 &= O_p \left(\frac{1}{N^\alpha R_N^2} \right) \times O_p(N) = o_p(1).
 \end{aligned}$$

Hence, from (A.9) and (A.2) we deduce that

$$\begin{aligned}
 \frac{(R_N^2 - 1) \sum_{t=1}^N x_t^2}{R_N^2 N^\alpha} &= \frac{R_N^2 x_N^2}{R_N^2 N^\alpha} \{1 + o_p(1)\} \\
 &= \left(\frac{x_N}{R_N^N N^{\alpha/2}} \right)^2 \{1 + o_p(1)\} \\
 &= \left\{ \frac{x_0}{N^{\alpha/2}} + \frac{1}{N^{\alpha/2}} \sum_{j=1}^N R_N^{-j} u_{xj} + \frac{N^{\alpha/2}}{c} \mu \right\}^2 \\
 &\quad \times \{1 + o_p(1)\} \\
 &\Rightarrow \left(\frac{\sigma_{xx}}{(2c)^{1/2}} U_x + D \right)^2. \tag{A.10}
 \end{aligned}$$

(iii) Combining the results (A.8) and (A.10), we have

$$\begin{aligned}
 \frac{(R_N^N N^\alpha)^{-1} \sum_{t=1}^N x_t u_{0t}}{\frac{(R_N^2 - 1) \sum_{t=1}^N x_t^2}{R_N^2 N^\alpha}} &\sim \frac{(R_N^N N^\alpha)^{-1} \sum_{t=1}^N x_t u_{0t}}{\frac{2c}{R_N^2 N^{2\alpha}} \sum_{t=1}^N x_t^2} \\
 &\Rightarrow \frac{\frac{\sigma_{00}}{(2c)^{1/2}} U_0 \left(D + \frac{\sigma_{xx}}{(2c)^{1/2}} U_x \right)}{\left(\frac{\sigma_{xx}}{(2c)^{1/2}} U_x + D \right)^2} \\
 &= \frac{\frac{\sigma_{00}}{(2c)^{1/2}} U_0}{\frac{\sigma_{xx}}{(2c)^{1/2}} U_x + D} = \frac{\sigma_{00} U_0}{\sigma_{xx} U_x + (2c)^{1/2} D}.
 \end{aligned}$$

Therefore,

$$R_N^N N^\alpha (\hat{A} - A) = \frac{(R_N^N N^\alpha)^{-1} \sum_{t=1}^N x_t u_{0t}}{(R_N^N N^{2\alpha})^{-1} \sum_{t=1}^N x_t^2} \Rightarrow 2c \frac{\sigma_{00} U_0}{\sigma_{xx} U_x + (2c)^{1/2} D},$$

giving the stated result. ■

A.2. Proof of Corollary 2.2

Proof. The proof follows from Theorem 2.1 by noting the mappings

$$\begin{aligned}
 \sigma_{00}^2 &\mapsto \sigma_{00}^2, \sigma_{xx}^2 \mapsto 1, R_N \mapsto a_h = e^{-\kappa h}, X^* \mapsto \frac{x_0}{\sigma_{xx}}, \\
 \mu &\mapsto \frac{\mu\kappa}{\sigma_{xx}} h^{1/2}, \mu^* \mapsto \frac{\mu\kappa}{\sigma_{xx}}, \\
 D_N &\mapsto D_h = \tilde{x}_0 h^{1/2} - \frac{h^{-1/2} \tilde{g}_h}{\kappa} \rightarrow D^* = \frac{x_0}{\sigma_{xx}} - \frac{\mu}{\sigma_{xx}},
 \end{aligned}$$

with $h = 1/N^\alpha$. It follows that

$$a_h^{-N} h \sum_{t=1}^N \tilde{x}_{th} u_{0,th} \Rightarrow \frac{\sigma_{00}}{-2\kappa} U_0 (U_x + (-2\kappa)^{1/2} D^*), \tag{A.11}$$

$$a_h^{-2N} h^2 \sum_{t=1}^N \tilde{x}_{th}^2 \Rightarrow \left(\frac{1}{-2\kappa} \right)^2 (U_x + (-2\kappa)^{1/2} D^*)^2, \tag{A.12}$$

and hence

$$\begin{aligned} \frac{a_h^N}{\sqrt{h}} (\widehat{\beta} - \beta) &\Rightarrow (-2\kappa) \frac{\frac{\sigma_{00}}{\sigma_{xx}} U_0}{U_x + (-2\kappa)^{1/2} D^*} \\ &= (-2\kappa) \frac{\sigma_{00} U_0}{\sigma_{xx} U_x + (-2\kappa)^{1/2} (x_0 - \mu)}. \quad \blacksquare \end{aligned}$$

Appendix B. Supplementary data

Supplementary material related to this article can be found online at <http://dx.doi.org/10.1016/j.jeconom.2017.08.016>.

References

- Aït-Sahalia, Y., Mykland, P.A., Zhang, L., 2005. How often to sample a continuous-time process in the presence of market microstructure noise. *Rev. Financ. Stud.* 18 (2), 351–416.
- Andrews, D.W.K., Guggenberger, P., 2012. Asymptotics for LS, GLS, and feasible GLS statistics in an AR(1) model with conditional heteroskedasticity. *J. Econometrics* 169 (2), 196–210.
- Bandi, F.M., Russell, J.R., 2006. Separating microstructure noise from volatility. *J. Financ. Econom.* 79 (3), 655–692.
- Boswijk, H.P., 2001. Block Local to Unity and Continuous Record Asymptotics. Tinbergen Institute Discussion Paper, No. 01-078/4.
- Gouriéroux, C., Zakoian, J.M., 2017. Local explosion modelling by non-causal process. *J. R. Stat. Soc. Ser. B Stat. Methodol.* 79 (3), 737–756.
- Hannan, E.J., 1970. *Multiple Time Series*. Wiley, New York.
- Hansen, P.R., Lunde, A., 2006. Realized variance and market microstructure noise. *J. Bus. Econom. Statist.* 24 (127–161).
- Magdalinos, T., Phillips, P.C.B., 2009. Limit theory for cointegrated systems with moderately integrated and moderately explosive regressors. *Econometric Theory* (482–526).
- Marsaglia, G., 1965. Ratios of normal variables and ratios of sums of uniform variables. *J. Amer. Statist. Assoc.* 60 (193–204).
- Phillips, P.C.B., 1972. The structural estimation of a stochastic differential equation system. *Econometrica* 40 (1021–1041).
- Phillips, P.C.B., 1991. Error correction and long run equilibria in continuous time. *Econometrica* 59 (967–980).
- Phillips, P.C.B., Magdalinos, T., 2007. Limit theory for moderate deviations from a unit root. *J. Econometrics* 136 (1), 115–130.
- Phillips, P.C.B., Magdalinos, T., 2009. Unit root and cointegrating limit theory when initialization is in the infinite past. *Econometric Theory* 25 (06), 1682–1715.
- Phillips, P.C.B., Magdalinos, T., 2013. Inconsistent VAR regression with common explosive roots. *Econometric Theory* 29 (4), 808–837.
- Phillips, P.C., Moon, H.R., Xiao, Z., 2001. How to estimate autoregressive roots near unity. *Econometric Theory* 17 (1), 29–69.
- Phillips, P.C.B., Shi, S., Yu, J., 2015a. Testing for multiple bubbles: Historical episodes of exuberance and collapse in the S&P 500. *Internat. Econom. Rev.* 56 (4), 1043–1078.
- Phillips, P.C.B., Shi, S., Yu, J., 2015b. Testing for multiple bubbles: Limit theory of dating algorithms. *Internat. Econom. Rev.* 52, 201–226.
- Phillips, P.C.B., Wu, Y., Yu, J., 2011. Explosive behavior in the 1990s NASDAQ: When did exuberance escalate asset values? *Internat. Econom. Rev.* 52 (201–226).
- Phillips, P.C.B., Yu, J., 2006. Comments on ‘Realized Variance and Market Microstructure Noise’ by P. R. Hansen and A. Lunde. *J. Bus. Econom. Statist.* 24, 202–208.
- Phillips, P.C.B., Yu, J., 2011. Dating the timeline of financial bubbles during the subprime crisis. *Q. Econ.* 2, 455–491.
- Wang, X., Yu, J., 2016. Double asymptotics for an explosive continuous time model. *J. Econometrics* 193, 35–53.
- Zhang, L., Mykland, P.A., Aït-Sahalia, Y., 2005. A tale of two time scales: Determining integrated volatility with noisy high-frequency data. *J. Amer. Statist. Assoc.* 100 (472), 1394–1411.